

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00626 A1

(51) International Patent Classification⁷: C07D 223/16,
A61K 31/55, A61P 13/00, C07D 401/06, 243/12, 495/04

Richard, Jeremy [GB/GB]; 208 Finchampstead Road,
Wokingham, Berkshire RG40 3HB (GB). **SEMPLE,**
Graeme [GB/SE]; AstraZeneca R & D Molndal, S-431 83
Molndal (SE).

(21) International Application Number: PCT/GB01/02737

(22) International Filing Date: 21 June 2001 (21.06.2001)

(74) Agent: **GEERING, Keith, Edwin**; Reddie & Grose, 16
Theobalds Road, London WC1X 8PL (GB).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0015601.8 26 June 2000 (26.06.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **FER-**
RING B.V. [NL/NL]; Polaris Avenue 144, NL-2132 JX
Hoofddorp (NL).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and



(75) Inventors/Applicants (*for US only*): **ASHWORTH,**
Doreen, Mary [GB/GB]; 10 The Glades, Locks Heath,
Southampton, Hampshire SO31 6UY (GB). **PITT,**
Gary, Robert, William [GB/GB]; 18 Kingfishers, Ship-
ton-Bellinger, Tidworth, Hampshire SP9 7US (GB).
HUDSON, Peter [GB/GB]; 12 Totteahale Close, North
Baddesley, Southampton, Hampshire SO52 9NQ (GB).
YEA, Christopher, Martyn [GB/GB]; 11 Gaston Gar-
dens, Romsey, Hampshire SO51 7RZ (GB). **FRANKLIN,**

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.


(54) Title: FUSED AZEPINE DERIVATIVES AND THEIR USE AS ANTIDIURETIC AGENTS



(57) Abstract: Compounds according to general formulae (1 and 2), wherein G¹ is an azepine derivative and G² is a group according to general formulae (9 - 11) are new. Compounds according to the invention are vasopressin V₂ receptor agonists. Pharmaceutical compositions of the compounds are useful as antidiuretic agents.

WO 02/00626 A1

In fact, such an agent is already in use in human therapy. Desmopressin (otherwise [1-desamino, D-Arg⁸]vasopressin, Minirin™, DDAVP™, Octostim™) is a peptide analogue of vasopressin which is selectively an agonist at the V₂ receptor. It is used in the treatment of central diabetes insipidus, which is a condition that results from defective secretion of vasopressin. It is also employed in the control of nocturnal enuresis and may also be of use in the control of nocturia. However, desmopressin is not an ideal agent in all respects. Even the best current syntheses of the agent are lengthy, and desmopressin is not amenable to the most convenient of purification techniques such as crystallisation. Consequently, desmopressin is relatively expensive. It has a very low oral bioavailability, and there is some variability in this parameter.




Overall then, there is a recognised need for a selective vasopressin V₂ receptor agonist that is easy to prepare and purify, and that has a high and predictable oral bioavailability. Such properties are most likely to be obtained with a non-peptide compound. Examples of such compounds are disclosed by Ogawa *et al.* in International Patent Application PCT/JP96/03652 (WO97/22591), by Failli *et al.* in PCT/US98/15487 (WO99/06403), PCT/US00/00885 (WO00/46224), and PCT/US00/00358 (WO00/46227), by Dusza *et al.* in PCT/US98/15495 (WO99/06409), and by Steffan and Failli in PCT/US00/00886 (WO00/46225), and PCT/US00/00658 (WO00/46228). However the compounds disclosed in these documents are not ideal drug candidates. For example, some have only moderate selectivity for the V₂ receptor and many have only very limited oral bioavailability, probably because they are poorly soluble in aqueous media. The present invention provides compounds that show a better combination of properties.

The anti-diuretic action of desmopressin results in a decrease in the osmolarity of the blood, and this has been shown to be useful in the treatment and prophylaxis of sickle-cell disease. Besides its antidiuretic actions, desmopressin is used to increase the

concentration in the blood of the coagulation proteins known as Faktor VIII and von Willebrand factor. In the clinical context, this makes desmopressin useful in the treatment of haemophilia A and von Willebrand's disease. Desmopressin has also been reported to show effects in the central nervous system. For example, it has been reported to be effective in the treatment of Tourette's disease and to be useful in the management of cocaine addiction. Similar applications would be open to the non-peptide agonists of the present invention.

SUMMARY OF THE INVENTION

The present invention relates to a series of compounds according to general formulae 1 and 2, and to salts and tautomers thereof, that are non-peptide agonists of vasopressin and which are selective for the V₂ receptor subtype.




wherein:


W is either N or C-R⁴;

R¹ – R⁴ are independently selected from H, F, Cl, Br, alkyl, CF₃, phenyl, OH, O-alkyl, NH₂, NH-alkyl, N(alkyl)₂, NO₂ and CN, or R² and R³ together can be -CH=CH-CH=CH-;


G¹ is a bicyclic or tricyclic fused azepine derivative selected from general formulae 3 to 8,




3




4




5



6



7



8

in which A^1 , A^4 , A^7 and A^{10} are each independently selected from CH_2 , O and NR^5 ;

A^2 , A^3 , A^9 , A^{11} , A^{13} , A^{14} and A^{15} are each independently selected from CH and N;

either A^5 is a covalent bond and A^8 is S, or A^5 is $\text{N}=\text{CH}$ and A^6 is a covalent bond;

A^8 and A^{12} are each independently selected from NH, N- CH_3 and S;

A^{16} and A^{17} are both CH_2 , or one of A^{16} and A^{17} is CH_2 and the other is selected from $\text{CH}(\text{OH})$, CF_2 , O, SO_a and NR^5 ;

R^5 is selected from H, alkyl and $(\text{CH}_2)_b\text{R}^6$;

R^6 is selected from phenyl, pyridyl, OH, O-alkyl, NH_2 , NH-alkyl, $\text{N}(\text{alkyl})_2$, NO_2 , CO_2H and CN;


a is 0, 1 or 2;

b is 1, 2, 3 or 4;


Y is CH or N;

Z is $\text{CH}=\text{CH}$ or S; and


G^2 is a group selected from general formulae 9 to 11,



9



10



11

in which Ar is selected from phenyl, pyridyl, naphthyl and mono- or polysubstituted phenyl or pyridyl wherein the substituents are selected from F, Cl, Br, alkyl, OH, O-alkyl, NH₂, NH-alkyl, N(alkyl)₂, NO₂ and CN;

D is a covalent bond or NH;

E¹ and E² are both H, OMe or F, or one of E¹ and E² is OH, O-alkyl, OBn, OPh, OAc, F, Cl, Br, N₃, NH₂, NHBn or NHAc and the other is H, or E¹ and E² together are =O, -O(CH₂)_gO- or -S(CH₂)_gS-;

F¹ and F² are both H, or together are =O or =S;

L is selected from OH, O-alkyl, NH₂, NH-alkyl and NR⁹R¹⁰;

R⁷ is selected from H, alkyl, alkenyl and COR⁸;

R⁸ is selected from OH, O-alkyl, NH₂, NH-alkyl, N(alkyl)₂, pyrrolidinyl and piperidinyl;

R⁹ and R¹⁰ are both alkyl, or together are -(CH₂)_h- or -(CH₂)₂O(CH₂)₂;

V is O, N-CN or S;

c is 0 or 1;

d is 0 or 1;

e is 0 or 1;

f is 0, 1, 2, 3 or 4;

g is 2 or 3; and


h is 3, 4 or 5,

provided that d and e are not both 0.

The invention further comprises pharmaceutical compositions incorporating these vasopressin agonists, which compositions are particularly useful in the treatment of central diabetes insipidus, nocturnal enuresis and nocturia.







DESCRIPTION OF THE INVENTION

In a first aspect, the present invention comprises novel 4-(aminomethyl)benzamide and 6-(aminomethyl)nicotinamide derivatives according to general formulae 1 and 2.



In general formula 1, W represents either a nitrogen atom (N) or a substituted carbon atom (C-R⁴). The substituents R¹ – R⁴ are each independently selected from hydrogen (H), fluorine (F), chlorine (Cl) and bromine (Br) atoms, and alkyl, trifluoromethyl (CF₃), phenyl (Ph), hydroxyl (OH), alkoxy (O-alkyl), primary amino (NH₂), monoalkylamino (NH-alkyl), dialkylamino (N(alkyl)₂), nitro (NO₂) and cyano (CN) groups. Alternatively, R² and R³ together can be -CH=CH-CH=CH- such that together with the ring to which they are attached they form a naphthalene, isoquinoline or isoquinolin-3-one fused ring system. The relationship between the two general formulae above is clear when one considers the compound of general formula 1 in which W is nitrogen and R¹ is hydroxyl. The resulting 2-hydroxypyridine can also exist as its 2-pyridone tautomer. In this tautomeric form the nitrogen atom is able to carry a substituent equivalent to R⁴, and such a compound is represented by general formula 2.

The group G¹ is a bicyclic or tricyclic fused azepine derivative selected from general formulae 3 to 8. It is joined to the carbonyl group of the parent molecule (1 or 2) through the nitrogen atom of the azepine ring common to all of 3 to 8, so as to form an amide bond.




In these formulae, A¹, A⁴, A⁷ and A¹⁰ each represent an oxygen atom (-O-) or a methylene (-CH₂-) or substituted imino (-NR⁵-) group. A², A³, A⁹, A¹¹, A¹³, A¹⁴ and A¹⁵ each represent a methine group (=CH-) or a nitrogen atom (=N-). Where two or more of these occur in the same group, each is independent of the others. Thus, for example, in formula 3, A² and A³ may both be nitrogen, both methine, or one may be methine and the other nitrogen. A⁵ and A⁶ are chosen together such that either A⁵ is a covalent bond and A⁶ is a sulphur atom (-S-), to give a thiophene ring, or A⁵ is a group -N=CH- and A⁶ is a covalent bond to give a pyridine ring. A⁸ and A¹² each represent an imino group (-NH-), an N-methyl imino group (-NCH₃-) or a sulphur atom (-S-). A¹⁶ and A¹⁷ may both represent a methylene group (-CH₂-) or one of A¹⁶ and A¹⁷ may represent a methylene group while the other represents a hydroxymethylene group (-CH(OH)-), a difluoromethylene group (-CF₂-), a substituted imino group (-NR⁵-), an oxygen atom (-O-), or an optionally oxidised sulphur atom (-SO_a-), where a is zero, 1 or 2.

The group R⁵ represents a hydrogen atom (H), an alkyl group, or a group -(CH₂)_bR⁶, where b is 1, 2, 3 or 4. The group R⁶ represents a group selected from phenyl, pyridyl, hydroxy (-OH), alkoxy (-O-alkyl), primary amino (-NH₂), mono- and dialkylamino (-NH-alkyl and N(alkyl)₂), nitro (-NO₂), carboxy (-CO₂H) and cyano (-CN) groups.

Y represents either a methine group (=CH-) or a nitrogen atom (=N-). Z represents either a sulphur atom (-S-) or a group -CH=CH-.

The group G^2 is selected from general formulae 9 to 11.



In these formulae, V represents a divalent residue selected from oxygen (=O) and sulphur (=S) atoms and a cyanimide (=N-CN) group.

In general formula 9, Ar represents an aromatic group selected from phenyl, pyridyl, naphthyl and mono- or polysubstituted phenyl and pyridyl groups, wherein the substituents are selected from fluorine (F), chlorine (Cl) and bromine (Br) atoms and alkyl, hydroxy (-OH), alkoxy (-O-alkyl), primary amino (-NH₂), mono- and dialkylamino (-NH-alkyl and N(alkyl)₂), nitro (-NO₂), carboxy (-CO₂H) and cyano (-CN) groups.

The values of c, d and e are independently zero or 1, provided that d and e are not both zero.

In general formula 10, D represents a covalent bond or an imino group (-NH-). The group R⁷ represents a hydrogen atom (H), an alkyl or alkenyl group, or a group -COR⁸, in which R⁸ represents a hydroxy (-OH), alkoxy (-O-alkyl), primary amino (-NH₂) or mono- or dialkylamino (-NH-alkyl and N(alkyl)₂) group, or a cyclic amino group selected from pyrrolidiny (-N(CH₂)₄) and piperidiny (-N(CH₂)₅). The value of f is zero, 1, 2, 3 or 4.

In general formula 11, E¹ and E² represent either two monovalent atoms or groups, which may be the same or different, or together they represent a divalent atom or group. When E¹ and E² represent monovalent atoms or groups, these may both simultaneously be hydrogen (H) or fluorine (F) atoms or methoxy (-OMe) groups, or one may be a fluorine (F), chlorine (Cl) or bromine (Br) atom, or a hydroxy (-OH), alkoxy (-O-alkyl), benzyloxy (-OBn), phenoxy (-OPh), acetoxy (-OAc), azido (-N₃), primary amino (-NH₂), benzylamino (-NH-Bn) or acetamido (-NHAc) group and the other is a hydrogen atom (H). When E¹ and E² together represent a divalent atom or group, this may be an oxygen atom (=O) or an α,ω-dioxa- or dithiapolyethylene group (-O(CH₂)_gO- or -S(CH₂)_gS-), in which the value of g is either 2 or 3.

F¹ and F² may both represent a hydrogen (H) atom. Alternatively, they may together represent an oxygen (=O) or sulphur (=S) atom. L represents a group selected from hydroxy (-OH), alkoxy (-O-alkyl), primary amino (-NH₂) and monoalkylamino (-NH-alkyl) groups and -NR⁹R¹⁰, wherein either R⁹ and R¹⁰ each represent alkyl groups which may be the same or different, or together they represent a polymethylene group -(CH₂)_h- in which h can be 3, 4 or 5, or -(CH₂)₂O(CH₂)₂-.

As used herein, the term "alkyl" includes saturated hydrocarbon residues, including linear, branched and cyclic groups, with up to six carbon atoms. Examples of alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl, isobutyl, *tert*-butyl, neopentyl and cyclohexyl groups.

The term "alkenyl" includes mono-unsaturated hydrocarbon residues, including linear, branched and cyclic groups, of between two and six carbon atoms. Examples of alkenyl groups include, but are not limited to, vinyl, 1-propenyl, allyl, 2-methyl-2-propenyl, 2-butenyl, 3-cyclopentenyl and 2,3-dimethyl-2-butenyl groups.

Certain compounds within the scope of the present invention may exist as tautomers. For example, when W is nitrogen and R¹ or R² is a hydroxy group, or when Ar is pyridyl further substituted by a hydroxy group, the resulting hydroxypyridine can exist as the pyridone tautomer. All such tautomers are considered to be within the scope of the present invention.

Certain compounds of general formula 1 are capable of forming salts with acids or bases. For example, compounds containing one or more basic nitrogen atoms can form addition salts with mineral and organic acids such as hydrochloric acid, sulphuric acid, phosphoric acid, acetic acid, trifluoroacetic acid, methanesulphonic acid, citric acid and benzoic acid. Compounds containing acidic groups can form salts with bases. Examples of such salts include the sodium, potassium, calcium, triethylammonium and tetraethylammonium salts. Furthermore, compounds that have both acidic and basic groups can form internal salts (zwitterions). Insofar as these salts are pharmaceutically acceptable, they are included within the scope of the invention.

A preferred embodiment of the invention is a compound according to general formula 1. More preferred is a compound according to general formula 1 in which W is C-R⁴. Even

more preferred is such a compound in which at least one of $R^1 - R^4$ is other than hydrogen. Most preferred is a compound in which one of $R^1 - R^4$ is methyl, chlorine or fluorine and the other three are hydrogen.

Another preferred embodiment of the invention is a compound according to general formula 2. More preferred is a compound according to general formula 2 in which R^2 and R^3 are both hydrogen.

Another preferred embodiment of the invention is a compound according to general formulae 1 or 2 in which G^1 is a group according to any of general formulae 3 to 7. More preferred is a compound in which Y is CH. Even more preferred is a compound in which Z is $-\text{CH}=\text{CH}-$ so as to complete a benzenoid ring. Alternatively, Z may be S to complete a thiophene ring. When Y is N it is particularly preferred that Z be $-\text{CH}=\text{CH}-$ so as to complete a pyridine ring.


Within the foregoing preferred embodiment, more preferred compounds are those wherein G^1 is a group according to general formula 3, particularly those wherein A^1 is CH_2 and both A^2 and A^3 are CH, and compounds wherein G^1 is a group according to general formula 6, particularly those wherein A^{11} is CH and A^{12} is S.

Another preferred embodiment of the invention is a compound according to general formulae 1 or 2 in which G^1 is a group according to general formula 8. More preferred is a compound in which one of A^{16} and A^{17} is CH_2 . Even more preferred is a compound in which both A^{16} and A^{17} are CH_2 .

Another preferred embodiment of the invention is a compound according to general formulae 1 or 2 in which G^2 is a group according to general formula 9. More preferred are those compounds wherein Ar is mono- or polysubstituted phenyl. Even more preferred are phenyl groups with at least two halogen (fluorine or chlorine) substituents. Most preferably, Ar is 2,6-difluorophenyl.

Another preferred embodiment of the invention is a compound according to general formulae 1 or 2 in which G^2 is a group according to general formula 10. More preferred are those compounds wherein R^7 is COR^8 . Most preferred are those compounds wherein R^8 is N(alkyl)_2 .

Another preferred embodiment of the invention is a compound according to general formulae 1 or 2 in which G^2 is a group according to general formula 11. More preferred are those compounds wherein F^1 and F^2 together are =O. Also preferred are those compounds wherein both E^1 and E^2 are H, or one is H and the other is O-alkyl. For those compounds wherein one of E^1 and E^2 is H and the other is O-alkyl, it is preferred that the stereochemistry at the CE^1E^2 centre be of the *R* absolute configuration. It is further preferred that the stereochemistry adjacent to the ring nitrogen atom be of the *S* absolute configuration. These configurations are illustrated below.



To the extent that the features of the foregoing preferred embodiments are independent of each other they may be combined in embodiments that are more preferred. Thus, highly preferred embodiments of the invention are those compounds that combine the preferred options for W and $R^1 - R^4$ with the preferred options for G^1 and G^2 .

A most preferred embodiment of the invention is a compound selected from the following.

1-(4-[3-(2-Chloro-6-fluorophenyl)ureidomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1*H*-1-benzazepine

1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5-(3-pyridyl)methyl-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepine

1-(3-Chloro-4-[3-(2-chloro-6-fluorophenyl)ureidomethyl]benzoyl)-5-ethyl-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepine

4-(3-Chloro-4-[3-(2,6-difluorophenyl)ureidomethyl]benzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine

1-(3-Chloro-4-(3-(methyloxycarbonyl)propanoylaminomethyl)benzoyl)-2,3,4,5-tetrahydro-1-benzazepine

1-(2-Methyl-4-(5-(3-pyridylmethyl)-2,3,4,5-tetrahydro-1,5-benzodiazepin-1-ylcarbonyl)benzyl)-3-(methyloxycarbonylmethyl)urea

1-(2-Methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide

(4*R*)-4-Hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide

(4*R*)-1-(3-Chloro-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide

(4*R*)-1-(2-Chloro-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide

(4*R*)-4-Benzyloxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide

(4*R*)-4-Methoxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide

(4*R*)-4-Methoxy-1-(3-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide

(4*R*)-1-(2-Chloro-4-(5,6,7,8-tetrahydro-4*H*-thieno[3,2-*b*]azepin-4-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide

(4*R*)-1-(4-(10,11-Dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-yl carbonyl)-2-methylbenzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide

(4*R*)-1-(2-Chloro-4-(10,11-Dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide

(4*R*)-1-(4-(10,11-Dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)-2-methylbenzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylthioamide

Within this set of compounds, two which show an optimal balance of properties are 1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide and (4*R*)-4-hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide.

The present invention further comprises pharmaceutical compositions that include at least one compound according to the foregoing description as an active constituent. The composition may also include a second pharmacological agent such as a spasmolytic or a potassium channel blocker, these agents being known in the art to ameliorate bladder dysfunction. Preferably, the composition includes only one active constituent. The composition will include excipients selected from binding agents, bulking agents, dispersants, solvents, stabilising agents and the like, such excipients being generally known in the art.

The excipients used will depend on the intended nature of the formulation, which will, in turn, depend on the intended route of administration. Administration may be oral, transmucosal (such as sublingual, buccal, intranasal, vaginal and rectal), transdermal or by injection (such as subcutaneous, intramuscular and intravenous). Oral administration is generally preferred. For oral administration, the formulation will be a tablet or capsule. Other formulations include dry powders, solutions, suspensions, suppositories and the like.

In a further aspect, the present invention is a method of treating or controlling certain human physiological dysfunctions. This method comprises the administration to the person in need of such treatment of an effective amount of a pharmaceutical composition, which composition contains a compound according to the foregoing description as an active constituent. The compounds act to reduce urine output, and so the method of the invention can be applied to all conditions in which elevated urine output is a contributory factor. The compounds also increase the production of the blood coagulation proteins known as Factor VIII and von Willebrand factor, and so the treatment of bleeding disorders can be undertaken.

In a preferred embodiment, the condition treated is central diabetes insipidus. This is a condition caused by an inability of the body to produce and secrete physiologically active vasopressin, with the result that water re-uptake is greatly reduced and large volumes of urine are produced.

In another preferred embodiment, the condition treated is nocturnal enuresis. This is defined as bladder emptying while the individual is sleeping. It is a condition that mainly affects children and a number of factors may be involved in its etiology.

In another preferred embodiment, the condition treated is nocturia. This is defined as production of sufficient urine during the night to require the individual to wake and empty his (or her) bladder. Again, this condition may be the result of a number of factors.

In another preferred embodiment, the condition treated is incontinence. This condition is characterised, in part, by reduced bladder capacity and control such that involuntary urination occurs unless the bladder is emptied frequently. Incontinence has been divided into two conditions, stress incontinence and urge incontinence. A number of etiological factors are thought to be involved. Treatment according to the invention is particularly useful for delaying the need for bladder emptying ("voiding postponement") in order to allow the incontinent subject a dry period of a few hours (such as up to four hours). Such voiding postponement may also be useful for the non-incontinent population, for example for people obliged to remain in meetings for extended periods.


In another preferred embodiment, the condition treated is haemophilia A or von Willebrand's disease. This is a condition in which Factor VIII or von Willebrand factor production is reduced and the individual suffers from prolonged bleeding.

In another preferred embodiment, the composition is administered prior to surgery (including dental surgery) to increase the coagulability of the blood and so reduce peri-operative blood loss.

The administration of the compositions of the present invention will generally be under the control of a physician. The physician will determine the amount of composition to be administered and the dosing schedule, taking into account the patient's physical condition and the therapeutic goals. For an adult diabetes insipidus patient, a typical dose might

be between 50mg and 1g of the active compound per day, taken as a single tablet or as up to four tablets throughout the day. For routes of administration other than the oral route, the amount of compound will be reduced, since non-oral routes tend to be more efficient in terms of delivering therapeutic agents into the systemic circulation. For the treatment of von Willebrand's disease and haemophilia A, the amount of compound may need to be higher than for the treatment of diabetes insipidus.

The compounds of the present invention can be prepared using methods generally known in the art. The compounds of general formulae 1 and 2 can be considered to be composed of three linked fragments, G^1 , G^2 and the central aromatic moiety (which will be referred to here as the "core"). Reagents corresponding to the three fragments will generally be prepared separately and then combined at a late stage in the synthesis.




Some instances of the various groups and substituents might be incompatible with this assembly and so will require the use of protecting groups. The use of protecting groups is well known in the art (see for example "Protective Groups in Organic Synthesis", T.W. Greene, Wiley-Interscience, 1981). Particular groups that may require protection are amines (protected as amides or carbamates), alcohols (protected as esters or ethers) and carboxylic acids (protected as esters). For the purposes of this discussion, it will be assumed that such protecting groups as are necessary are in place.

The three fragments can be combined according to two strategies to give the compounds of formulae 1 and 2. In the first, the fragments corresponding to G^1 and the core are linked to give a fragment corresponding to core- G^1 , which is then combined with fragment G^2 . In the second, the fragments corresponding to the core and G^2 are linked to give a fragment corresponding to G^2 -core, which is then combined with fragment G^1 . The chemistry involved in the condensation of fragment G^1 with the core fragment,

and that involved in the condensation of the core fragment with fragment G², will be the same whichever strategy is followed.


Formation of fragment core-G¹




The synthesis of this fragment requires the formation of an amide bond between the two moieties. Reactions of this type are well known in the art. Most conveniently, an acid chloride corresponding to the core fragment may be allowed to react with the free secondary amino group of the G¹ azepine ring. Such a reaction generally is performed in an aprotic solvent such as dichloromethane or dimethylformamide at or slightly below room temperature. A tertiary amine base such as triethylamine or dimethylaminopyridine is usually added. Alternatively, the carboxylic acid corresponding to the core fragment may be condensed with the secondary amino group using one of the many reagents that have been developed for the formation of amide bonds in the field of peptide chemistry. Examples of such reagents include DCC (dicyclohexylcarbodiimide), BOP ((benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate), PyBOP[®] ((benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate), PyBroP[®] (bromotripyrrolidinophosphonium hexafluorophosphate) and HBTU (O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate). Other reagents are also known. The details of the synthetic method will depend on the particular reagent selected, but will generally involve the use of an aprotic solvent and a tertiary amine base, as described above. Either the reagent is added to a mixture of the carboxylic acid and the azepine, or the carboxylic acid and the reagent are premixed to form a reactive intermediate (which is not isolated) to which is added the azepine.

Formation of fragment G²-core

Depending on the nature of G², the G²-core bond can be part of an amide or thioamide, a sulphonamide, a urea or thiourea, a sulphonylurea or sulphonylthiourea, or a cyanoamidine, cyanoguanidine or sulphonylcyanoguanidine. The chemistry involved in the preparation of the G²-core bond will be different for each of these.


(i.a) Amides {G² = 10, D = covalent bond, V = O}

These compounds can be formed by the reaction of a carboxylic acid or acid chloride corresponding to fragment G² with the primary amino group of the core fragment. Conditions for the reaction will generally be similar to those described for the formation of the core-G¹ bond, except that the primary amine is more reactive than the azepine nitrogen and so lower temperatures and shorter reaction times may be used.

(i.b) Thioamides {G² = 10, D = covalent bond, V = S}


These compounds can be formed by the reaction of a suitable thiocarbonyl compound such as a dithioester (RCS₂R') with the primary amine in a manner analogous to that described for the corresponding amides above. Alternatively, they may be prepared from the corresponding amides (V = O) by reaction with Lawesson's reagent.

(ii) *Sulphonamides* { $G^2 = 9$, $d = 1$, $e = \text{zero}$ }



These compounds are generally prepared by the reaction of the sulphonyl chloride corresponding to the G^2 fragment with the primary amine of the core fragment. The reaction is generally performed under conditions similar to those described above for the reaction of a carboxylic acid chloride with the primary amine that gives the amides.

(iii.a) *Ureas* { $G^2 = 9$, $d = \text{zero}$, $e = 1$, $V = O$; $G^2 = 10$, $D = NH$, $V = O$; $G^2 = 11$, $V = O$ }




These compounds can be prepared by the reaction of an amine with an isocyanate or an equivalent thereof. Due to the symmetry of the urea functional group, there is the possibility to choose which component acts as the amine and which as the isocyanate. Most simply, when G^2 is a group according to 9 or 10, the corresponding isocyanate is readily accessible. It can conveniently be reacted with the primary amine of the core

fragment in an aprotic solvent without the need for additional reagents. When G^2 is a group according to 11, the isocyanate is not available, and the carbamoyl chloride can be used in its place. The carbamoyl chloride is generally prepared immediately prior to use by treating the corresponding secondary amine with phosgene or an equivalent reagent such as diphogene or triphogene. Alternatively, the use of carbonyl diimidazole leads to the formation of a carbamoyl imidazole derivative that can be used in place of the carbamoyl chloride. The reaction of the carbamoyl chloride with the primary amine generally requires the addition of a tertiary amine base to neutralise the hydrogen chloride formed.


In some cases, it may be preferable to treat the primary amine corresponding to the core fragment with phosgene (or carbonyl diimidazole) to form an isocyanate that can subsequently be reacted with the primary or secondary amine corresponding to the G^2 fragment.

(iii.b) Thioureas { $G^2 = 9$, $d = \text{zero}$, $e = 1$, $V = S$; $G^2 = 10$, $D = NH$, $V = S$; $G^2 = 11$, $V = S$ }



These compounds can be prepared by methods analogous to those described above for the ureas, simply by using the corresponding isothiocyanate and thiophosgene compounds.


(iv.a) Sulphonylureas { $G^2 = 9$, $d = 1$, $e = 1$, $V = O$ }



These compounds can be prepared by the reaction of the primary amine corresponding to the core fragment with an appropriate sulphonyl isocyanate. The reaction conditions


are similar to those described above for the reaction of an amine with an isocyanate to prepare the ureas.

(iv.b) Sulphonylthioureas $\{G^2 = 9, d = 1, e = 1, V = S\}$




These compounds can be prepared analogously to the sulphonylureas by the reaction of the primary amine corresponding to the core fragment with an appropriate sulphonyl isothiocyanate.

(v.a) Cyanoamidines $\{G^2 = 10, D = \text{covalent bond}, V = N-CN\}$




These compounds can be prepared by the reaction of the primary amine of the core fragment with an *N*-cyanothioamide or an *N*-cyanothioimide corresponding to the G^2 fragment.

(v.b) Cyanoguanidines $\{G^2 = 9, d = \text{zero}, e = 1, V = N-CN; G^2 = 10, D = NH, V = N-CN; G^2 = 11, V = N-CN\}$



These compounds can be prepared by the reaction of the primary amine of the core fragment with a cyanothiourea corresponding to the G^2 fragment in the presence of a carbodiimide.

(v.c) Sulphonylcyanoguanidines ($G^2 = 9$, $d = 1$, $e = 1$, $V = N-CN$)



These compounds can be prepared in an analogous manner by the reaction of the primary amine of the core fragment with an *N*-sulphonyl-*N'*-cyanothiourea corresponding to the G^2 fragment in the presence of a carbodiimide.

The reagents corresponding to the fragments are commercially available, or they can be prepared by methods described in the literature. Particularly relevant leading references include the following.

Synthesis of fused azepine derivatives for G^1 :

Aranapakam *et al.*, Bioorg. Med. Chem. Lett. 1993, 1733; Artico *et al.*, Farmaco. Ed. Sci. 24, 1969, 276; Artico *et al.*, Farmaco. Ed. Sci. 32, 1977, 339; Chakrabarti *et al.*, J. Med. Chem. 23, 1980, 878; Chakrabarti *et al.*, J. Med. Chem. 23, 1980, 884; Chakrabarti *et al.*, J. Med. Chem. 32, 1989, 2573; Chimirri *et al.*, Heterocycles 36, 1993, 601; Grunewald *et al.*, J. Med. Chem. 39, 1996, 3539; Klunder *et al.*, J. Med. Chem. 35, 1992, 1887; Liegeois *et al.*, J. Med. Chem. 37, 1994, 519; Olagbemiro *et al.*, J. Het. Chem. 19, 1982, 1501; Wright *et al.*, J. Med. Chem. 23, 1980, 462; Yamamoto *et al.*, Tet. Lett. 24, 1983, 4711; and International patent application, publication number WO99/06403.

Synthesis of amidine transfer reagents for G^2 , $V = N-CN$

Mestres *et al.*, Synthesis, 1980, 755; Petersen *et al.*, J. Med. Chem. 21, 1978, 773; and Cord, J. Chem. Soc., 1948, 1620.

Synthesis of proline derivatives for $G^2 =$ group according to 11

Dugave *et al.*, Tet. Lett. 39, 1998, 1169; Petrillo *et al.*, J. Med. Chem. 31, 1988, 1148; and Smith *et al.*, J. Med. Chem. 31, 1988, 875.

The foregoing general description is further illustrated below with a number of non-limiting examples.

EXAMPLES


Abbreviations

The following abbreviations have been used.

AIBN	Azo- <i>bis</i> -(isobutyronitrile)
BOC	<i>tert</i> -Butyloxycarbonyl
(BOC) ₂ O	Di- <i>tert</i> -butyl dicarbonate
DMF	Dimethylformamide
EtOAc	Ethyl acetate
IPA	Isopropanol
M.S.	Mass spectrometry
NBS	N-Bromosuccinimide
pet. ether	petroleum ether, fraction boiling at 60-80°C
THF	Tetrahydrofuran
WSCDI	Water-soluble carbodiimide

Preparation of Intermediates


Reagents corresponding to fragments G¹ and G² were commercially available or prepared according to the published procedures except where detailed in the specific Examples. Reagents corresponding to the core fragment were prepared as detailed below.

Example A**4-(tert-Butyloxycarbonylaminoethyl)-3-chlorobenzoic acid****A1. Methyl 4-bromomethyl-3-chlorobenzoate**


To a solution of methyl 3-chloro-4-methylbenzoate (5.0g, 27.1mmol) in carbon tetrachloride (50ml) were added NBS (5.8g, 32.0mmol) and AIBN (0.442g, 2.70mmol). The mixture was stirred at reflux for 18h. The mixture was allowed to cool to room temperature and then concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 0:100 to 5:95); yield 5.96g (84%).

A2. 4-(tert-Butyloxycarbonylaminoethyl)-3-chlorobenzoic acid


To a saturated solution of ammonia in ethanol (170ml) was added methyl 4-bromomethyl-3-chlorobenzoate from Example A1 (5.5g, 20.9mmol). The mixture was stirred at room temperature for 1hr and then concentrated *in vacuo*. The residue was triturated with diethyl ether and the resultant white crystals were filtered off and washed with more diethyl ether. To a solution of this solid in water (100ml) were added solutions of (BOC)₂O (5.0g, 23.0mmol) in dioxan (100ml) and sodium hydroxide (1.86g, 46.0mmol) in water (100ml). The mixture was stirred at room temperature for 18h and then concentrated *in vacuo*. The aqueous residue was acidified with citric acid and extracted with chloroform/IPA. The organic layer was washed with water, dried over MgSO₄, and concentrated *in vacuo* to give a white solid; yield 2.8g (67%).

Example B**4-(tert-Butyloxycarbonylaminomethyl)-3-nitrobenzoic acid**


4-Bromomethyl-3-nitrobenzoic acid (4.75g, 18.2mmol) was reacted following the method of Example A2 to give a yellow solid; yield 2.6g (49%).

Example C**4-Cyano-3-methylbenzoic acid**

To a solution of 4-bromo-2-methylbenzonitrile (2.0g, 10.2mmol) in THF (100ml) at -78°C under a nitrogen atmosphere was added dropwise a 2.5M solution of *n*-butyl lithium (4.48ml, 11.2mmol). The mixture was stirred at -78°C for 1h and then poured onto solid carbon dioxide (5g) in THF (50ml). The mixture was allowed to warm to room temperature. Water was added (200ml) and the mixture was extracted with diethyl ether (3 times). The aqueous layer was acidified by addition of concentrated HCl and extracted with chloroform (3 times). The combined chloroform extracts were washed with water, dried over MgSO_4 , and concentrated *in vacuo* to give a white solid; yield 1.2g (73%).

Example D**4-Cyano-2-methylbenzoic acid**

4-Bromo-3-methylbenzonitrile (2.0g, 10.2mmol) was reacted following the method of Example C to give a yellow solid which was triturated with hexane and filtered off; yield 0.96g (59%).

Example E**4-(*tert*-Butyloxycarbonylaminoethyl)-2-fluorobenzoic acid****E1. 2-Fluoro-4-methylbenzoic acid**

4-Bromo-3-fluorotoluene (8.33g, 44.07mmol) was reacted following the method of Example C to give a white solid; 4.89g (72%).

E2. Methyl 2-fluoro-4-methylbenzoate

To a solution of 2-fluoro-4-methylbenzoic acid from Example E1 (6.04g, 39.18mmol) in toluene (80ml) was added thionyl chloride (65ml, 89.11mmol). The mixture was heated at reflux for 2.5h, cooled and concentrated *in vacuo*. The residue was dissolved in

dichloromethane (50ml) and methanol (50ml) was added. The mixture was stirred at room temperature for 2.5h and then concentrated *in vacuo*. The residue was dissolved in dichloromethane (100ml), washed with saturated sodium bicarbonate solution and brine, dried over MgSO_4 , and concentrated *in vacuo* to give a tan solid; yield 5.07g (77%).

E3. Methyl 4-bromomethyl-2-fluorobenzoate


Methyl 2-fluoro-4-methylbenzoate from Example E2 (5.07g, 30.16mmol) was reacted following the method of Example of A1. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 20:80); yield 5.9g (80%).

E4. 4-(*tert*-Butyloxycarbonylaminomethyl)-2-fluorobenzoic acid

Methyl 4-bromomethyl-2-fluorobenzoate from Example E3 (5.9g, 24.13mmol) was reacted following the method of Example A2. The product was recrystallised from dioxan/pet. ether to give white crystals; yield 2.46g (38%).

Example F

6-(*tert*-Butyloxycarbonylaminomethyl)-2-chloronicotinic acid



F1. Methyl 2-chloro-6-methylnicotinate

To a suspension of 2-chloro-6-methylnicotinic acid (5.3g, 30.8mmol) in dichloromethane (100ml) at 0°C were added DMF (1ml) and oxalyl chloride (3.2ml, 36.9mmol). The mixture was allowed to warm to room temperature and stirred for 5h. The solvents were removed *in vacuo* and the residue was dissolved in dichloromethane (50ml) and methanol (50ml). The mixture was stirred at room temperature for 18h and then

concentrated *in vacuo*. The residue was dissolved in chloroform, washed with saturated sodium bicarbonate solution and brine, dried over MgSO_4 , and concentrated *in vacuo* to give a brown oil; yield 5.70g (100%).

F2. Methyl 6-bromomethyl-2-chloronicotinate


Methyl 2-chloro-6-methylnicotinate from Example F1 (5.70g, 30.8mmol) was reacted following the method of Example of A1. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 20:80); yield 4.8g (58%).

F3. Methyl 6-(*tert*-butoxycarbonylaminomethyl)-2-chloronicotinate

Methyl 6-bromomethyl-2-chloronicotinate from Example F2 (4.8g, 18.0mmol) was reacted following the method of Example of A2 to give an off white solid; yield 1.45g (28%).

Example G

6-(*tert*-Butyloxycarbonylaminomethyl)nicotinic acid



G1. Methyl 6-(bromomethyl)nicotinate

Methyl 6-methylnicotinate (5.0g, 33.0mmol) was reacted following the method of Example A1. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 20:80); yield 3.7g (49%).

G2. Methyl 6-(azidomethyl)nicotinate


To a solution of methyl 6-(bromomethyl)nicotinate from Example G1 (2.0g, 8.60mmol) in DMF (15ml) was added sodium azide (0.84g, 12.9mmol). The mixture was stirred at room temperature for 18h. EtOAc (100ml) was added and the mixture was washed with water (3 times), dried over MgSO_4 , and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 20:80) to give a yellow gum; yield 1.55g (93%).

G3. Methyl 6-(tert-butyloxycarbonylaminomethyl)nicotinate

To a degassed solution of methyl 6-(azidomethyl)nicotinate from Example G2 (1.6g, 8.30mmol) in methanol (50ml) was added 10% palladium-on-carbon (0.15g). Hydrogen gas was bubbled through the mixture for 2h at room temperature. The catalyst was removed by filtration through a pad of celite and the filtrate was evaporated *in vacuo*. The residue was dissolved in dichloromethane and cooled to 0°C. To this solution were added triethylamine (1.67g, 16.0mmol) and $(\text{BOC})_2\text{O}$ (2.17g, 9.96mmol). The mixture was allowed to warm to room temperature and stirred for 18h, then concentrated *in vacuo*. The residue was dissolved in EtOAc and washed with water, dried over MgSO_4 , and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50) to give a yellow solid; yield 1.57g (71%).

G4. 6-(tert-Butyloxycarbonylaminomethyl)nicotinic acid

To a solution of methyl 6-(tert-butyloxycarbonylaminomethyl)nicotinate from Example G3 (1.56g, 5.84mmol) in THF (20ml) and water (5ml) was added lithium hydroxide monohydrate (0.37g, 8.76mmol). The mixture was stirred at room temperature for 18h and then concentrated *in vacuo*. The aqueous residue was acidified by addition of 1M citric acid solution and extracted with chloroform/IPA (3 times). The combined organic extracts were washed with brine, dried over MgSO_4 , and evaporated *in vacuo* to give a white solid; yield 1.38g (94%).

Example H**4/5-Bromo-6-(tert-butyloxycarbonylaminomethyl)-1-methyl-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid****H1. Methyl 1,6-dimethyl-2-oxo-1,2-dihydropyridine-3-carboxylate**

To a solution of 3-hydroxy-6-methylnicotinic acid (10g, 65.0mmol) in DMF (100ml) at 0°C was added sodium hydride (4.83g, 60% dispersion, 140mmol). The mixture was stirred at 0°C for 1.5h, then methyl iodide (12.4ml, 195mmol) was added and the mixture was allowed to warm to room temperature, stirring for a further 18h. The mixture was partitioned between water and EtOAc and the aqueous layer acidified to pH 5. The layers were separated and the organic layer was washed with brine, dried over MgSO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant dichloromethane/methanol 95:5) to give a white solid. This was recrystallised from methanol and the filtrate was evaporated *in vacuo* to give the desired product; yield 6.1g (52%).

H2. Methyl 4/5-bromo-6-bromomethyl-1-methyl-2-oxo-1,2-dihydropyridine-3-carboxylate


Methyl 1,6-dimethyl-2-oxo-1,2-dihydropyridine-3-carboxylate of Example H1 (6.0g, 33.0mmol) was reacted following the method of Example of A1. The product was purified by flash chromatography on silica (eluant dichloromethane/methanol 95:5); yield 5.2g (46%).

H3. 4/5-Bromo-6-(tert-butyloxycarbonylaminomethyl)-1-methyl-2-oxo-1,2-dihydropyridine-3-carboxylic acid

Methyl 4/5-bromo-6-bromomethyl-1-methyl-2-oxo-1,2-dihydropyridine-3-carboxylate of Example H2 (5.2g, 14.8mmol) was reacted following the method of Example A2 to give a brown gum; yield 1.3g (24%).

Example I

4-Cyano-3,5-dimethylbenzoic acid



I1. 4-Bromo-2,6-dimethylbenzonitrile

4-Bromo-2,6-dimethylaniline (4.49g, 22.4mmol) was taken up in water (25ml) and concentrated hydrochloric acid (8.0ml) was added. The mixture was sonicated to form a fine suspension and then cooled to 0°C. A solution of sodium nitrite (1.67g, 24.2mmol) in water (5ml) was then added dropwise so as to maintain the temperature of the reaction between 0-5°C. The mixture was stirred at 0-5°C for 1/2h and then neutralised by addition of solid sodium carbonate. The resulting solution was then added portionwise to a solution of copper cyanide (2.42g, 27.0mmol) and potassium cyanide (3.65g, 56.1mmol) in water (25ml) at 70°C. The mixture was stirred at 70°C for 1/2h, allowed to cool and then extracted with toluene (2 times). The combined extracts were washed with water and brine, dried over MgSO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 5:95) to give an orange solid; yield 3.2g (68%).


12. 4-Cyano-3,5-dimethylbenzoic acid

4-Bromo-2,6-dimethylbenzonitrile from Example I1 (3.20g, 15.2mmol) was reacted following the method of Example C to give a tan solid; yield 1.5g (56%).

Reagents corresponding to fragments A, B and C were combined to give the specific Examples as detailed below.

Example 1

1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine



1A. 1-(4-Cyanobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

To a solution of 2,3,4,5-tetrahydro-1H-1-benzazepine (1.05g, 7.14mmol) in dichloromethane (40ml) were added 4-cyanobenzoic acid (1.26g, 8.57mmol), triethylamine (1.00g, 7.14mmol), 4-(dimethylamino)pyridine (0.87g, 7.14mmol) and WSCDI (2.86g, 14.28mmol). The mixture was stirred at reflux for 18h, cooled and evaporated *in vacuo*. The residue was partitioned between EtOAc and 1M KHSO₄. The organic layer was washed with saturated sodium bicarbonate solution and brine, dried over MgSO₄, and concentrated *in vacuo* to give a white solid; yield 1.50g (76%).


1B. 1-(4-(Aminomethyl)benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

To a degassed solution of the cyanobenzoyl benzazepine from Example 1A (1.50g, 5.43mmol) in methanol (50ml) were added concentrated hydrochloric acid (1.4ml, 16.2mmol) and 10% palladium-on-carbon (1.15g). Hydrogen gas was bubbled through the mixture for 5h at room temperature. The catalyst was removed by filtration through a pad of celite and the filtrate was evaporated *in vacuo*. The residue was partitioned between EtOAc and water. The aqueous layer was basified by addition of saturated sodium bicarbonate solution and extracted with dichloromethane (2 times). The combined organic extracts were washed with brine, dried over MgSO₄, and concentrated *in vacuo* to give a white solid; yield 1.12g (74%).

1C. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine


To a solution of the amine from Example 1B (0.50g, 1.79mmol) in dichloromethane (20ml) were added triethylamine (0.27ml, 1.97mmol) and 2,6-difluorophenylisocyanate (0.31g, 1.97mmol). The mixture was stirred at room temperature for 2h and then evaporated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50) to give a white solid; yield 0.62g (80%).

M.S.: calc m/e=435.18; found [M+H]⁺= 436.

Example 2.**1-(4-[3-(2,6-Difluorophenyl)cyanoguanidinomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

To a solution of the amine from Example 1B (0.12g, 0.379mmol) in DMF (20ml) were added 1-(2,6-difluoro-phenyl)-3-cyano-thiourea (0.16g, 0.758mmol, prepared according to Atwal *et. al.*, Tetrahedron Lett., 30, p7313, 1989.), diisopropylethylamine (0.16ml, 0.947mmol) and WSCDI (0.087g, 0.455mmol). The mixture was stirred at room temperature for 72h and then evaporated *in vacuo*. The residue was partitioned between dichloromethane and 1M KHSO₄. The organic layer was washed with saturated sodium bicarbonate solution and brine, dried over MgSO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50-70:30) to give a white solid; yield 0.084g (48%).

M.S.: calc m/e=459.19; found [M+H]⁺= 460.0

Example 3.**1-(6-[3-(2,6-Difluorophenyl)ureidomethyl]nicotinoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine****3A. 1-[6-(*tert*-Butyloxycarbonylaminomethyl)nicotinoyl]-2,3,4,5-tetrahydro-1H-1-benzazepine**

The carboxylic acid from Example G4 (1.38g, 5.45mmol) was reacted with 2,3,4,5-tetrahydro-1H-1-benzazepine (0.80g, 5.50mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70-70:30); yield 1.14g (55%).

3B. 1-[6-(Aminomethyl)nicotinoyl]-2,3,4,5-tetrahydro-1H-1-benzazepine hydrochloride

The BOC amine from Example 3A (1.14g, 2.98mmol) was dissolved in 4N HCl/dioxan, stirred at room temperature for 1h and then evaporated *in vacuo*, azeotroping with toluene, to give an off white solid; yield 1.0g (quantitative).


3C. 1-(6-[3-(2,6-Difluorophenyl)ureidomethyl]nicotinoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The amine hydrochloride from Example 3B (0.070g, 0.220mmol) was reacted with 2,6-difluorophenylisocyanate (0.038g, 0.242mmol) according to the procedure in Example 1C. The product was purified by trituration with diethyl ether to give a white solid; yield 0.060g (63%).

M.S.: calc $m/e=436.47$; found $[M+H]^+=437.2$.

Example 4.

1-(3-Chloro-4-[3-(3-methoxyphenyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine



4A. 1-(4-[*tert*-Butyloxycarbonylaminomethyl]-3-chlorobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The carboxylic acid from Example A2 (1.0g, 3.50mmol) was reacted with 2,3,4,5-tetrahydro-1H-1-benzazepine (0.47g, 3.20mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70-40:60); yield 0.88g (66%).

4B. 1-(4-[Aminomethyl]-3-chlorobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine hydrochloride

The BOC amine from Example 4A (0.88g, 2.10mmol) was dissolved in 4N HCl/dioxan and stirred at room temperature for 1h, then evaporated *in vacuo*, azeotroping with toluene, to give a white solid; yield 0.70g (95%).


4C. 1-(3-Chloro-4-[3-(3-methoxyphenyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The amine hydrochloride from Example 4B (0.050g, 0.140mmol) was reacted with 3-methoxyphenylisocyanate (0.021g, 0.140mmol) according to the procedure in Example 1C. The product was purified by trituration with diethyl ether to give a white solid; yield 0.060g (93%).

M.S.: calc $m/e=463.17$; found $[M+H]^+=464.2$.


Example 5.

1-(3-Chloro-4-[3-(2-chlorophenyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine




The amine hydrochloride from Example 4B (0.050g, 0.140mmol) was reacted with 2-chlorophenylisocyanate (0.022g, 0.140mmol) according to the procedure in Example 1C. The product was purified by trituration with diethyl ether to give a white solid; yield 0.063g (98%).

M.S.: calc $m/e=467.12$; found $[M+H]^+$; $^{35}\text{Cl}=468.1$.

Example 6.**1-(3-Chloro-4-[3-(2,6-difluorophenyl)thioureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

The amine hydrochloride from Example 4B (0.075g, 0.214mmol) was reacted with 2,6-difluorophenylisothiocyanate (0.054g, 0.320mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70-45:55); yield 0.068g (66%).

M.S.: calc m/e=485.11; found $[M+H]^+$; ^{35}Cl = 486.2, $[M+H]^+$; ^{37}Cl = 488.1

Example 7.**1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-2-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

7A. 1-(4-Cyano-2-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The carboxylic acid from Example D (0.96g, 5.95mmol) was reacted with 2,3,4,5-tetrahydro-1H-1-benzazepine (0.80g, 5.44mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70); yield 0.59g (38%).


7B. 1-(4-[Aminomethyl]-2-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine hydrochloride

The cyanobenzoyl benzazepine from Example 7A (0.59g, 2.03mmol) was hydrogenated according to the procedure in Example 1B. The product was isolated as the HCl salt; yield 0.55g (82%).

7C. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-2-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The amine hydrochloride from Example 7B (0.050g, 0.151mmol) was reacted with 2,6-difluorophenylisocyanate (0.028g, 0.181mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50); yield 0.041g (62%).

M.S.: calc m/e=449.19; found $[M+H]^+$ = 450.1.

Example 8.**1-(3-Methyl-4-[3-(phenylsulfonyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine****8A. 1-(4-Cyano-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

The carboxylic acid from Example C (0.96g, 5.95mmol) was reacted with 2,3,4,5-tetrahydro-1H-1-benzazepine (0.80g, 5.44mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70); yield 1.10g (70%).


8B. 1-(4-[Aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine hydrochloride

The cyanobenzoyl benzazepine from Example 8A (1.10g, 3.79mmol) was hydrogenated according to the procedure in Example 1B. The product was isolated as the HCl salt; yield 1.23g (98%).

8C. 1-(3-Methyl-4-[3-(phenylsulfonyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine


The amine hydrochloride from Example 8B (0.050g, 0.151mmol) was reacted with phenylsulphonylisocyanate (0.028g, 0.151mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 80:20); yield 0.026g (22%).

M.S.: calc $m/e=477.17$; found $[M+H]^+=478.2$.

Example 9.**1-(3-Methyl-4-[3-(2-oxo-1,2-dihydropyrid-3-yl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**


To a suspension of 2-hydroxynicotinic acid (95mg, 0.68mmol) in dioxan (5ml) were added triethylamine (0.11ml, 0.771mmol) and diphenylphosphoryl azide (0.16ml, 0.725mmol). The mixture was stirred at reflux for 3h. The amine hydrochloride from Example 8B (0.15g, 0.453mmol) and triethylamine (0.095ml, 0.680mmol) were added and the mixture was stirred at reflux for a further 18h, cooled and evaporated *in vacuo*. The residue was partitioned between dichloromethane and 1M KHSO₄. The organic layer was washed with saturated sodium bicarbonate solution and brine, dried over MgSO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant methanol:dichloromethane 2:98-5:95) to give a white solid; yield 0.084g (43%).

M.S.: calc m/e=430.20; found [M+H]⁺= 431.1.

Example 10.**1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

The amine hydrochloride from Example 8B (0.050g, 0.151mmol) was reacted with 2,6-difluorophenylisocyanate (0.028g, 0.181mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50); yield 0.044g (65%).

M.S.: calc $m/e=449.19$; found $[M+H]^+=450.1$.

Example 11.**1-(3-Nitro-4-[2-nitrobenzylsulfonylaminoethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

11A. 1-(4-[*tert*-Butyloxycarbonylamino
methyl]-3-nitrobenzoyl)-2,3,4,5-tetrahydro-1*H*-1-
benzazepine

The carboxylic acid from Example B (0.911g, 3.08mmol) was reacted with 2,3,4,5-tetrahydro-1*H*-1-benzazepine (0.453g, 3.08mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50); yield 0.58g (43%).


11B. 1-(4-[Aminomethyl]-3-nitrobenzoyl)-2,3,4,5-tetrahydro-1*H*-1-benzazepine
hydrochloride

The BOC-aminomethylbenzoyl benzazepine from Example 11A (0.33g, 0.764mmol) was reacted according to the procedure in Example 4B. The product was isolated as the HCl salt; yield 0.27g (98%).

11C. 1-(3-Nitro-4-[2-nitrobenzylsulfonylamino
methyl]benzoyl)-2,3,4,5-tetrahydro-1*H*-1-
benzazepine

The amine hydrochloride from Example 11B (0.068g, 0.188mmol) was reacted with 2-nitrobenzylsulphonyl chloride (0.033g, 0.226mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 25:75-50:50); yield 0.010g (10%).

M.S.: calc m/e=524.14; found $[M+H]^+$ = 525.2.

Example 12.**1-(3-Amino-4-[3-(2,6-difluorophenyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine****12A. 1-(3-Amino-4-[tert-butyloxycarbonylaminomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

To a degassed solution of the nitrobenzoyl benzazepine from Example 11A (0.30g, 0.700mmol) in methanol (50ml) was added 10% palladium-on-carbon (0.10g). Hydrogen gas was bubbled through the mixture for 1.5h at room temperature. The catalyst was removed by filtration through a pad of celite and the filtrate was evaporated *in vacuo*; yield 0.254g (92%).

12B. 1-(3-Amino-4-[aminomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine dihydrochloride

The BOC-aminomethylbenzoyl benzazepine from Example 12A (0.14g, 0.354mmol) was reacted according to the procedure in Example 4B. The product was isolated as the diHCl salt; yield 0.098g (75%).

12C. 1-(3-Amino-4-[3-(2,6-difluorophenyl)ureidomethyl]benzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine


The amine hydrochloride from Example 12B (0.132g, 0.35mmol) was reacted with 2,6-difluorophenylisocyanate (0.055g, 0.35mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether

70:30) and then by preparative HPLC (gradient water:acetonitrile 80:20-20:80; 0.1% TFA). The HPLC fractions were freeze-dried to give a white solid; yield 0.027g (17%).

M.S.: calc $m/e=450.19$; found $[M+H]^+=451.2$.

Example 13.

1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-dimethylaminobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine



13A. 1-(4-[tert-Butyloxycarbonylaminomethyl]-3-dimethylaminobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

To an ice cold solution of the amine from Example 12A (0.16g, 0.40mmol) in 1% acetic acid/methanol (25ml) was added formaldehyde (37% solution in water, 0.050ml, 0.60mmol). The mixture was stirred at 0°C for 10min and then sodium borohydride (0.050g, 0.80mmol) was added. The mixture was allowed to warm to room temperature with stirring over 1h. and then evaporated *in vacuo*. The residue was partitioned between EtOAc and saturated sodium bicarbonate solution. The organic layer was washed with brine, dried over $MgSO_4$, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70-70:30) to give a white solid; yield 0.091g (56%).

13B. 1-(4-[Aminomethyl]-3-dimethylaminobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The BOC-aminomethylbenzoyl benzazepine from Example 13A (0.089g, 0.225mmol) was reacted according to the procedure in Example 4B. The product was isolated as the HCl salt; yield 0.075g (97%).


13C. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-dimethylaminobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The amine hydrochloride from Example 13B (0.075g, 0.20mmol) was reacted with 2,6-difluorophenylisocyanate (0.032g, 0.20mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 90:10); yield 0.044g (65%).

M.S.: calc m/e=478.22; found $[M+H]^+$ = 479.2.

Example 14.

1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-2-fluorobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine



14A. 1-(4-[tert-Butyloxycarbonylaminomethyl]-2-fluorobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The carboxylic acid from Example E4 (0.60g, 2.22mmol) was reacted with 2,3,4,5-tetrahydro-1H-1-benzazepine (0.28g, 1.89mmol) according to the procedure in Example

1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 40:60); yield 0.58g (77%).

14B. 1-(4-[Aminomethyl]-2-fluorobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The BOC-aminomethylbenzoyl benzazepine from Example 14A (0.58g, 1.42mmol) was reacted according to the procedure in Example 4B. The product was isolated as the HCl salt; yield 0.29g (60%).


14C. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-2-fluorobenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The amine hydrochloride from Example 14B (0.040g, 0.12mmol) was reacted with 2,6-difluorophenylisocyanate (0.020g, 0.13mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 40:60-100:0); yield 0.038g (70%).

M.S.: calc m/e=453.17; found $[M+H]^+$ = 454.1.

Example 15.

1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine



15A. 2,3,4,5-Tetrahydro-1H-1,5-benzodiazepine

To an ice cold solution of lithium aluminium hydride (4.68g, 123mmol) in dry THF (100ml), under a nitrogen atmosphere, was added dropwise a solution of 2,3,4,5-tetrahydro-1H-1,5-benzodiazepin-2-one (5.0g, 30.9mmol) in dry THF (50ml). The mixture was allowed to warm to room temperature and then heated at reflux for 2h. The mixture was then cooled to 0°C and a solution of aqueous ammonium hydroxide (10ml) in THF (60ml) was added dropwise. The resultant suspension was stirred for 1h and then filtered through a pad of celite. The filtrate was evaporated *in vacuo* to give a tan solid; yield 4.36g (95%).

15B. 1-(4-Cyano-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The carboxylic acid from Example C (0.65g, 4.03mmol) was reacted with 2,3,4,5-tetrahydro-1H-1,5-benzodiazepine from Example 15A (0.50g, 3.36mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50); yield 0.36g (37%).


15C. 1-(4-[Aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine hydrochloride

The cyanobenzoyl benzodiazepine from Example 15B (0.36g, 1.24mmol) was hydrogenated according to the procedure in Example 1B. The product was isolated as the HCl salt; yield 0.17g (40%).

15D. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The amine hydrochloride from Example 15C (0.170g, 0.46mmol) was reacted with 2,6-difluorophenylisocyanate (0.071g, 0.46mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 80:20); yield 0.089g (43%).

M.S.: calc m/e=450.19; found $[M+H]^+$ = 451.2.

Example 16.**1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5-(3-pyridyl)methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine****16A. 1-(3-Pyridyl)methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine**

To solution of 2,3,4,5-tetrahydro-1H-1,5-benzodiazepine from Example 15A (0.50g, 3.38mmol) in 1% acetic acid/methanol (25ml), at room temperature, was added pyridine-3-carboxaldehyde (0.35ml, 03.72mmol). The mixture was stirred at reflux for 18h and then allowed to cool to room temperature. Sodium borohydride (0.050g, 0.80mmol) was added. The mixture was stirred for 2h and then evaporated *in vacuo*. The residue was partitioned between EtOAc and saturated sodium bicarbonate solution. The organic layer was washed with brine, dried over MgSO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc) to give a white solid; yield 0.386g (40%).

16B. 1-(4-Cyano-3-methylbenzoyl)-5-(3-pyridyl)methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The carboxylic acid from Example C (0.31g, 1.93mmol) was reacted with 1-(3-pyridyl)methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine from Example 16A (0.39g,

1.61mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc); yield 0.28g (45%).


16C. 1-(4-Aminomethyl-3-methylbenzoyl)-5-(3-pyridyl)methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

To a solution of the nitrile from Example 16B (0.28g, 0.72mmol) in methanol (20ml) were added cobaltous chloride (0.338g, 1.42mmol) and sodium borohydride (0.27g, 7.20mmol). The mixture was stirred at room temperature for 1h and then saturated aqueous ammonium chloride solution (10ml) was added. The mixture was concentrated *in vacuo* and the aqueous residue was partitioned between diethyl ether and water. The aqueous layer was basified by addition of saturated sodium bicarbonate solution and extracted with chloroform (3 times). The combined organic extracts were washed with brine, dried over MgSO₄, and concentrated *in vacuo* to give a white solid; yield 0.20g (72%).

16D. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5-(3-pyridyl)methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The amine from Example 16C (0.065g, 0.168mmol) was reacted with 2,6-difluorophenylisocyanate (0.027g, 0.17mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc); yield 0.068g (75%).

M.S.: calc m/e=541.23; found [M+H]⁺= 542.2.

Example 17.**1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5-(2-hydroxyethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine****17A. Methyl (2-oxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepin-1-yl)acetate**

To a solution of 1,3,4,5-tetrahydro-benzo[b][1,4]diazepin-2-one (5.0g, 30.8mmol) in DMF (30ml), at -10°C, was added sodium hydride (1.35g, 60% dispersion, 33.9mmol). The mixture was stirred at -10°C for 15min, then methyl bromoacetate (2.92ml, 30.8mmol) was added. The mixture was stirred at -10°C for a further 1h and then concentrated *in vacuo*. The residue was taken up in EtOAc and washed with brine (3 times), dried over MgSO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc) to give a white solid; yield 7.08g (98%).

17B. 2-(2,3,4,5-Tetrahydro-1H-1,5-benzodiazepin-1-yl)ethanol

Methyl (2-oxo-[1,3,4,5-tetrahydro-benzo[b][1,4]diazepin-1-yl)-acetate from Example 17A (7.08g, 30.2mmol) was reduced with lithium aluminium hydride according to the procedure in Example 15A; yield 4.33g (75%).

17C. 1-(4-Cyano-3-methylbenzoyl)-5-(2-hydroxyethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

To a solution of the carboxylic acid from Example 1C (1.38g, 8.58mmol) in dichloromethane (50ml) was added thionyl chloride (3.33ml, 43.0mmol). The mixture was stirred at reflux for 2h and then evaporated *in vacuo*, azeotroping with toluene (2 times). The residue was dissolved in dichloromethane (50ml) and 2-(2,3,4,5-tetrahydro-1H-1,5-benzodiazepin-1-yl)ethanol from Example 17B (1.5g, 7.80mmol) was added. The mixture was stirred at room temperature for 18h and then evaporated *in vacuo*. The residue was partitioned between EtOAc and saturated sodium bicarbonate solution. The organic layer was washed with brine, dried over MgSO₄, and concentrated *in vacuo*. The residue was triturated with EtOAc and the resultant solid filtered off; yield 1.25g (48%).


17D. 1-(4-Aminomethyl-3-methylbenzoyl)-5-(2-hydroxyethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The cyanobenzoyl benzodiazepine from Example 17C (1.25g, 3.73mmol) was hydrogenated according to the procedure in Example 1B. The product was isolated as the free base; yield 0.94g (74%).

17E. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5-(2-hydroxyethyl)-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The amine from Example 17D (0.94g, 2.76mmol) was reacted with 2,6-difluorophenylisocyanate (0.47g, 3.04mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc); yield 0.068g (75%).

M.S.: calc m/e=494.21; found [M+H]⁺= 495.2.

Example 18.**1-(3-Chloro-4-[3-(2,6-difluorophenyl)ureidomethyl]benzoyl)-5-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine****18A. 1-Methyl-2-oxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine**

To a solution of 2,3,4,5-tetrahydro-1H-1,5-benzodiazepin-2-one (2.0g, 12.3mmol) in DMF (30ml), at -10°C, was added sodium hydride (0.54g, 60% dispersion, 13.6mmol). The mixture was stirred at -10°C for 15min, then methyl iodide (0.77ml, 12.3mmol) was added. The mixture was stirred at -10°C for a further 1h and then concentrated *in vacuo*. The residue was taken up in EtOAc and washed with brine (3 times), dried over MgSO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc) to give a white solid; yield 1.70g (78%).

18B. 1-Methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

1-Methyl-2-oxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine from Example 18A (1.7g, 9.65mmol) was reduced with lithium aluminium hydride according to the procedure in Example 15A; yield 1.34g (86%).

18C. 1-(4-[tert-Butyloxycarbonylaminoethyl]-3-chlorobenzoyl)-5-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The carboxylic acid from Example A2 (0.506g, 1.77mmol) was reacted with 1-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine from Example 18B (0.24g, 1.48mmol)

according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50); yield 0.30g (47%).

18D. 1-(4-Aminomethyl-3-chlorobenzoyl)-5-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The BOC-aminomethylbenzoyl benzazodiazepine from Example 18C (0.30g, 0.698mmol) was reacted according to the procedure in Example 4B. The product was isolated as the HCl salt; yield 0.25g (98%).


18E. 1-(3-Chloro-4-[3-(2,6-difluorophenyl)ureidomethyl]benzoyl)-5-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The amine hydrochloride from Example 18D (0.060g, 0.164mmol) was reacted with 2,6-difluorophenylisocyanate (0.021g, 0.164mmol) according to the procedure in Example 1C. The product was purified by trituration with diethyl ether to give a white solid; yield 0.058g (87%).

M.S.: calc m/e=484.15; found $[M+H]^+$; ^{35}Cl = 485.1.

Example 19.

1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-2-methylbenzoyl)-5-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine



19A. 1-(4-Cyano-2-methylbenzoyl)-5-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine hydrochloride

The carboxylic acid from Example D (0.50g, 3.10mmol) was reacted with 1-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine from Example 18B (0.46g, 2.80mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70-70:30); yield 0.27g (32%).


19B. 1-(4-Aminomethyl-2-methylbenzoyl)-5-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine hydrochloride

The cyanobenzoyl benzazepine from Example 19A (0.26g, 0.88mmol) was hydrogenated according to the procedure in Example 1B. The product was isolated as the HCl salt; yield 0.30g (99%).

19C. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-2-methylbenzoyl)-5-methyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The amine hydrochloride from Example 19B (0.060g, 0.17mmol) was reacted with 2,6-difluorophenylisocyanate (0.027g, 0.17mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 80:20); yield 0.070g (93%).

M.S.: calc m/e=464.20; found $[M+H]^+$ = 465.2.

Example 20.**1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3,5-dimethylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine****20A. 1-(4-Cyano-3,5-dimethylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

The carboxylic acid from Example 12 (0.49g, 2.80mmol) was reacted with 2,3,4,5-tetrahydro-1H-1-benzazepine (0.39g, 2.63mmol) according to the procedure in Example 17C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70); yield 0.66g (77%).


20B. 1-(4-Aminomethyl-3,5-dimethylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The nitrile from Example 20A (0.65g, 2.12mmol) was reduced according to the procedure in Example 16C; yield 0.42g (64%).

20C. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3,5-dimethylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The amine from Example 20B (0.070g, 0.23mmol) was reacted with 2,6-difluorophenylisocyanate (0.043g, 0.28mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 40:60); yield 0.033g (31%).

M.S.: calc m/e=463.21; found $[M+H]^+$ = 464.2.

Example 21.**1-(2-Chloro-6-[3-(2,6-difluorophenyl)ureidomethyl]nicotinoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine****21A. 1-(6-[*tert*-Butylaminomethyl]-2-chloronicotinoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

The carboxylic acid from Example F3 (0.50g, 1.74mmol) was reacted with 2,3,4,5-tetrahydro-1H-1-benzazepine (0.26g, 1.74mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 55:45); yield 0.038g (5%).

21B. 1-(6-Aminomethyl-2-chloronicotinoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine hydrochloride

The BOC-aminomethylnicotinoyl benzazepine from Example 21A (0.036g, 0.074mmol) was reacted according to the procedure in Example 4B. The product was isolated as the HCl salt; yield 0.026g (98%).

21C. 1-(2-Chloro-6-[3-(2,6-difluorophenyl)ureidomethyl]nicotinoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine


The amine hydrochloride from Example 21B (0.026g, 0.073mmol) was reacted with 2,6-difluorophenylisocyanate (0.014g, 0.08mmol) according to the procedure in Example

1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 90:10); yield 0.031g (90%).

M.S.: calc $m/e=470.13$; found $[M+H]^+$; $^{35}\text{Cl} = 471.1$.

Example 22.

1-(6-[3-(2,6-Difluorophenyl)ureidomethyl]-1-methyl-2-oxo-1,2-dihydropyridyl-3-carbonyl)-2,3,4,5-tetrahydro-1H-1-benzazepine



22A. 1-(4/5-Bromo-6-[tert-butyloxycarbonylaminomethyl]-1-methyl-2-oxo-1,2-dihydropyridyl-3-carbonyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The carboxylic acid from Example H3 (1.30g, 3.60mmol) was reacted with 2,3,4,5-tetrahydro-1H-1-benzazepine (0.53g, 3.60mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 60:40); yield 0.70g (40%).

22B. 1-(4/5-Bromo-6-[tert-butyloxycarbonylaminomethyl]-1-methyl-2-oxo-1,2-dihydropyridyl-3-carbonyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The benzazepine from Example 22A (0.60g, 1.23mmol) was hydrogenated according to the procedure in Example 12A; yield 0.50g (99%).

22C. 1-(6-Aminomethyl-1-methyl-2-oxo-1,2-dihydropyridyl-3-carbonyl)-2,3,4,5-tetrahydro-1H-1-benzazepine hydrochloride

The BOC-aminomethyl pyridone from Example 22B (0.50g, 1.22mmol) was reacted according to the procedure in Example 4B. The product was isolated as the HCl salt; yield 0.43g (99%).


22D. 1-(6-[3-(2,6-Difluorophenyl)ureidomethyl]-1-methyl-2-oxo-1,2-dihydropyridyl-3-carbonyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

The amine hydrochloride from Example 22C (0.050g, 0.144mmol) was reacted with 2,6-difluorophenylisocyanate (0.025g, 0.144mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:methanol 90:10); yield 0.064g (95%).

M.S.: calc m/e=466.18; found $[M+H]^+$ = 467.2.

Example 23.

1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5-ethyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine



23A. 1-Ethyl-2-oxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

2-Oxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine (1.95g, 11.96mmol) was reacted with ethyl iodide (1.4ml, 17.5mmol) according to the procedure in Example 18A. The product was purified by flash chromatography on silica (eluant EtOAc); yield 1.70g (75%).

23B. 1-Ethyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

1-Ethyl-2-oxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine from Example 23A (1.7g, 8.94mmol) was reduced with lithium aluminium hydride according to the procedure in Example 15A; yield 1.55g (98%).

23C. 1-(4-Cyano-3-methylbenzoyl)-5-ethyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The carboxylic acid from Example C (0.53g, 3.29mmol) was reacted with 1-ethyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine from Example 23B (0.514g, 2.92mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 60:40); yield 0.55g (59%).


23D. 1-(4-Aminomethyl-3-methylbenzoyl)-5-ethyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine hydrochloride

The nitrile from Example 23C (0.55g, 1.73mmol) was hydrogenated according to the procedure in Example 1B. The product was isolated as the HCl salt; yield 0.60g (96%).

23E. 1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5-ethyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepine

The amine hydrochloride from Example 23D (0.071g, 0.20mmol) was reacted with 2,6-difluorophenylisocyanate (0.038g, 0.25mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50-100:0); yield 0.044g (46%).

M.S.: calc m/e=478.22; found $[M+H]^+$ = 479.2.

Example 24.**5-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-6,7,8,9-tetrahydro-5H-pyrido[2,3-b]azepine****24A. 5-(4-Cyano-3-methylbenzoyl)-6,7,8,9-tetrahydro-5H-pyrido[2,3-b]azepine**

The carboxylic acid from Example C (0.36g, 2.26mmol) was reacted with 6,7,8,9-tetrahydro-5H-pyrido[2,3-b]azepine (0.33g, 2.23mmol) according to the procedure in Example 17C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 80:20); yield 0.47g (73%).

24B. 5-(4-Aminomethyl-3-methylbenzoyl)-6,7,8,9-tetrahydro-5H-pyrido[2,3-b]azepine

The cyanobenzoyl pyridoazepine from Example 24A (0.46g, 1.58mmol) was hydrogenated according to the procedure in Example 1B. The product was isolated as the free base; yield 0.28g (60%).


24C. 5-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-6,7,8,9-tetrahydro-5H-pyrido[2,3-b]azepine

The amine from Example 24B (0.071g, 0.20mmol) was reacted with 2,6-difluorophenylisocyanate (0.035g, 0.23mmol) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc); yield 0.020g (19%).

M.S.: calc $m/e=450.19$; found $[M+H]^+=451.2$.

Example 25.

5-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-1-oxo-1 λ^4 -2,3,4,5-tetrahydro-1,5-benzothiazepine



25A. 5-(4-Cyano-3-methylbenzoyl)-2,3,4,5-tetrahydro-1,5-benzothiazepine

The carboxylic acid from Example C (0.27g, 1.68mmol) was reacted with 2,3,4,5-tetrahydro-1,5-benzothiazepine (0.28g, 1.70mmol) according to the procedure in Example 1A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 60:40); yield 0.43g (84%).

25B. 5-(4-Aminomethyl-3-methylbenzoyl)-2,3,4,5-tetrahydro-1,5-benzothiazepine

The cyanobenzoyl benzothiazepine from Example 25A (0.43g, 1.40mmol) was hydrogenated according to the procedure in Example 1B. The product was isolated as the free base; yield 0.10g (29%).

25C. 5-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1,5-benzothiazepine

The amine from Example 25B (0.10g, 0.32mmol) was reacted with 2,6-difluorophenylisocyanate (0.061g, 0.39mmol) according to the procedure in Example

1C. The product was purified by trituration with diethyl ether to give a white solid; yield 0.112g (75%).


25D. 5-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-1-oxo-1 λ^4 -2,3,4,5-tetrahydro-1,5-benzothiazepine

To a suspension of the thiazepine from Example 25C (0.15g, 0.33mmol) in methanol (40ml), dichloromethane (10ml) and water (10ml) was added sodium periodate (0.21g, 0.99mmol). The mixture was stirred at room temperature for 70h and then filtered. The filtrate was evaporated *in vacuo* and the residue was purified by flash chromatography on silica (eluant EtOAc); yield 0.013g (8%).

M.S.: calc $m/e=483.14$; found $[M+H]^+=484.1$.

Example 26.

4-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-b]azepine



26A. 4-(4-Cyano-3-methylbenzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-b]azepine

The carboxylic acid from Example C (0.50g, 3.10mmol) was reacted with 5,6,7,8-tetrahydro-4H-thieno[3,2-b]azepine (0.45g, 2.95mmol) according to the procedure in Example 1A. The product was purified by recrystallisation from EtOAc:pet. ether; yield 0.48g (55%).

26B. 4-(4-Aminomethyl-3-methylbenzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-b]azepine

The nitrile from Example 26A (0.48g, 1.60mmol) was reduced according to the procedure in Example 16C; yield 0.16g (33%).


26C. 4-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-b]azepine

The amine from Example 26B (0.05g, 0.18mmol) was reacted with 2,6-difluorophenylisocyanate (0.027g, 0.18mmol) according to the procedure in Example 1C. The product was purified by trituration with diethyl ether to give a white solid; yield 0.052g (67%).

M.S.: calc m/e=455.15 found $[M+H]^+$ = 456.1.

Example 27.

4-(3-Methyl-4-[3-(2,3,5,6-tetrafluorophenyl)ureidomethyl]benzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-b]azepine




The amine from Example 26B (0.062g, 0.206mmol) was reacted with 2,3,5,6-tetrafluorophenylisocyanate (0.079g, 0.413mmol, prepared from the aniline according to the procedure of Kurita. K, *et. al.*, *J. Org. Chem.*, **41**, 1976, p2070.) according to the procedure in Example 1C. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50); yield 0.045g (44%).

M.S.: calc m/e=491.13 found $[M+H]^+$ = 492.1.

Example 28.

1-(4-[N-(4-Methoxy-4-oxobutanoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine



28A. 1-(4-Cyano-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

To a solution of 2,3,4,5-tetrahydro-1H-1-benzazepine (0.80g, 5.44mmol) in dichloromethane (40ml) were added 4-cyano-3-methylbenzoic acid from example C (0.96g, 5.95mmol), triethylamine (0.76g, 5.44mmol), 4-(dimethylamino)pyridine (0.66g, 5.44mmol) and WSCDI (2.17g, 10.88mmol). The mixture was stirred at reflux for 18h, cooled and evaporated *in vacuo*. The residue was partitioned between EtOAc and 1M KHSO₄. The organic layer was washed with saturated sodium bicarbonate solution and brine, dried over MgSO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70); yield 1.10g (70%).

28B. 1-(4-[Aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine hydrochloride

To a degassed solution of the cyanobenzoyl benzazepine from Example 28A (1.10g, 3.79mmol) in methanol (40ml) were added concentrated hydrochloric acid (0.98ml, 11.3mmol) and 10% palladium-on-carbon (0.80g). Hydrogen gas was bubbled through the mixture for 5h at room temperature. The catalyst was removed by filtration through a

pad of celite and the filtrate was evaporated *in vacuo* to give the product as the HCl salt; yield 1.23g (98%).


28C. 1-(4-[N-(4-Methoxy-4-oxobutanoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine

To a solution of the amine from Example 28B (0.10g, 0.30mmol) in dichloromethane (10ml) were added triethylamine (0.061ml, 0.90mmol) and 3-carbomethoxy propionyl chloride (0.046g, 0.30mmol). The mixture was stirred at room temperature for 18h and then washed with 1M KHSO₄ (3 times), water and brine, dried over Na₂SO₄, and concentrated *in vacuo* to give a white solid; yield 0.10g (81%).

M.S.: calc m/e=408; found [M+H]⁺= 409.


Example 29

1-(4-[N-(2-Methoxy-2-oxoethanoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine




The amine hydrochloride from Example 28B (0.10g, 0.30mmol) was reacted with methyl oxalyl chloride (0.037g, 0.30mmol) according to the procedure in Example 28C to give a white solid; yield 0.085g (76%).

M.S.: calc m/e=380; found [M+H]⁺= 381.

Example 30**1-(4-[N-(2-Hydroxy-2-oxoethanoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

To a solution of the methyl ester from Example 29 (0.045g, 0.118mmol) in THF (10ml) and water (5ml) was added lithium hydroxide monohydrate (0.010g, 0.23mmol). The mixture was stirred at room temperature for 2h, acidified to pH1 by addition of 1M HCl and extracted with EtOAc (3 times). The combined organic extracts were washed with brine, dried over Na₂SO₄, and concentrated *in vacuo* to give a white solid; yield 0.034g (76%).

M.S.: calc m/e=366; found [M+H]⁺= 367.


Example 31**1-(4-[N-(5-Methoxy-5-oxopentanoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

The amine hydrochloride from Example 28B (0.10g, 0.30mmol) was reacted with methyl 4-(chloroformyl) butyrate (0.050g, 0.30mmol) according to the procedure in Example 1C to give a white solid; yield 0.061g (48%).

M.S.: calc m/e=422; found $[M+H]^+$ = 423.


Example 32

1-(4-[N-(2-Ethoxy-2-oxoethylcarbamoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine




To a solution of the amine from Example 28B (0.10g, 0.30mmol) in dichloromethane (10ml) were added triethylamine (0.061ml, 0.90mmol) and ethyl isocyanatoacetate (0.059g, 0.45mmol). The mixture was stirred at room temperature for 18h and then washed with 1M KHSO₄ (3 times), water and brine, dried over Na₂SO₄, and concentrated *in vacuo* to give a white solid; yield 0.10g (81%).

M.S.: calc m/e=423; found $[M+H]^+$ = 424.

Example 33**1-(4-[N-(Carboxymethylcarbamoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

To a solution of the ethyl ester from Example 32 (0.050g, 0.10mmol) in THF (20ml) and water (5ml) was added lithium hydroxide monohydrate (0.020g, 0.45mmol). The mixture was stirred at room temperature for 4h. The mixture was concentrated *in vacuo* and the residue diluted with water then washed with diethyl ether. The aqueous layer was acidified to pH 1 by addition of 1M HCl and extracted with EtOAc (3 times). The combined organic extracts were washed with brine, dried over Na₂SO₄, and concentrated *in vacuo* to give a white solid; yield 0.046g (99%).

M.S.: calc m/e=395; found [M+H]⁺= 396.


Example 34**1-(4-[N-(2-Methylamino-2-oxoethyl)carbamoyl]aminomethyl)-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

To a solution of the carboxylic acid from Example 33 (0.10g, 0.25mmol) in dichloromethane (25ml) was added DIEA (0.221ml, 1.26mmol) and PyBroP (0.129g, 0.278mmol). The mixture was stirred at room temperature for 10min and then methylamine hydrochloride (0.085g, 1.26mmol) was added. Stirring was continued for a further 3h. The mixture was then washed with 1M KHSO₄ (3 times), saturated sodium bicarbonate solution (3 times) and brine, dried over Na₂SO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant dichloromethane:methanol 96:4) to give a white solid; yield 0.018g (17%).

M.S.: calc m/e=408; found [M+H]⁺= 409.


Example 35

1-(4-[N-(2-Dimethylamino-2-oxoethylcarbamoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine




The carboxylic acid from Example 33 (0.07g, 0.18mmol) was reacted with dimethylamine hydrochloride (0.072g, 0.88mmol) according to the procedure in Example 7. The product was purified by flash chromatography on silica (eluant chloroform:methanol:acetic acid 98:1:1) to give a white solid; yield 0.08g (11%).

M.S.: calc m/e=422; found [M+H]⁺= 423.

Example 36**1-(4-[N-(2-Methoxy-2-oxoethylcarbamoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**


To a solution of the carboxylic acid from Example 33 (0.080g, 0.20mmol) under a nitrogen atmosphere in dichloromethane (25ml) at 0°C were added DMF (20ml) and oxalyl chloride (31mg, 0.24mmol). The mixture was stirred at 0°C to room temperature for 2h and then concentrated *in vacuo*. The residue was dissolved in methanol (4ml) and dichloromethane (16ml) and the mixture stirred at room temperature for 16h. The mixture was then washed with 1M KHSO₄ (3 times), saturated sodium bicarbonate solution (3 times) and brine, dried over Na₂SO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant dichloromethane:methanol 96:4) to give a white solid; yield 0.049g (60%).

M.S.: calc m/e=409; found [M+H]⁺= 410.

Example 37**1-(4-[N-(2-Amino-2-oxoethylcarbamoyl)aminomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-1-benzazepine**

To a solution of the carboxylic acid from Example 33 (0.10g, 0.25mmol) in dichloromethane (20ml) were added hydroxybenzotriazole (34mg, 0.25mmol) and WSCDI (51mg, 0.25mmol). The mixture was stirred at room temperature for 10min. Ammonia 880 (0.5ml) was then added and stirring continued for a further 16h. The mixture was concentrated *in vacuo* and the residue purified by flash chromatography on silica (eluant ethyl acetate) to give a white solid; yield 0.008g (8%).

M.S.: calc $m/e=394$; found $[M+H]^+=395$.

Example 38**4-(4-[N-(4-Methoxy-4-oxobutanoyl)aminomethyl]-3-chlorobenzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-b]azepine**

38A. 4-(4-[N-(*tert*-Butyloxycarbonyl)aminomethyl]-3-chlorobenzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-*b*]azepine hydrochloride

The carboxylic acid from Example A2 (0.60g, 2.10mmol) was reacted with 5,6,7,8-tetrahydro-4H-thieno[3,2-*b*]azepine (0.28g, 1.80mmol) according to the procedure in example 28A. The product was purified by flash chromatography on silica (eluant EtOAc:pet. ether 40:60) to give a yellow solid.


38B. 4-(4-[Aminomethyl]-3-chlorobenzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-*b*]azepine hydrochloride

The BOC amine from Example 38A was dissolved in 4N HCl/dioxan (30 ml). The mixture was stirred at room temperature for 40min then concentrated *in vacuo* to leave a tan solid; yield 0.41g (63%, for 2 steps).

38C. 4-(4-[N-(4-Methoxy-4-oxobutanoyl)aminomethyl]-3-chlorobenzoyl)-5,6,7,8-tetrahydro-4H-thieno[3,2-*b*]azepine

To a solution of the amine from Example 38B (0.032g, 0.08mmol) in dichloromethane (10ml) were added triethylamine (0.025ml, 0.18mmol) and 3-carbomethoxypropionyl chloride (0.014g, 0.08mmol). The mixture was stirred at room temperature for 18h and then washed with 1M KHSO₄ (3 times), water and brine, dried over Na₂SO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 50:50-90:10); yield 0.022g (56%).

M.S.: calc m/e=434; found [M+H]⁺ ³⁵Cl = 435.

Example 39**1-(2-Methyl-4-((2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-N,N-dimethylamide****39A. 2-Methyl-4-((2,3,4,5-tetrahydro-1H-benzo[b]azepine)-1-carbonyl)-benzonitrile.**

To a solution of 2,3,4,5-tetrahydro-1H-benzo[b]azepine (0.80g, 5.44mmol) in dichloromethane (50ml) were added 4-cyano-3-methylbenzoic acid (0.96g, 5.95mmol), triethylamine (0.60g, 5.95mmol), 4-(dimethylamino)pyridine (0.73g, 5.95mmol) and WSCDI (1.24g, 6.48mmol). The mixture was stirred at reflux for 18h, cooled and evaporated *in vacuo*. The residue was partitioned between EtOAc and 1M KHSO₄. The organic layer was washed with saturated sodium bicarbonate solution and brine, dried over MgSO₄, and concentrated *in vacuo*. The crude material was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70); yield 1.10g (70%).

39B. 1-(4-(Aminomethyl)-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-benzo[b]azepine hydrochloride.

To a degassed solution of the cyanobenzazepine of Example 39A (1.10g, 3.79mmol) in methanol (50ml) were added concentrated hydrochloric acid (0.98ml, 11.3mmol) and 10% palladium on carbon (0.80g). Hydrogen gas was bubbled through the mixture for 5h at room temperature. The catalyst was removed by filtering through a pad of celite and the filtrate was evaporated; yield 1.23g (98%).


39C. 1-(2-Methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-N,N-dimethylamide

To a solution of the amine of Example 39B (0.10g, 0.302mmol) in DMF (10ml), under a nitrogen atmosphere, were added *N,N*-diisopropylethylamine (43mg, 0.332mmol) and carbonyl diimidazole (0.074g, 0.453mmol). The mixture was stirred at room temperature for 40 minutes. A solution of proline-*N,N*-dimethylamide (0.107g, 0.756mmol) in DMF (1ml) was added. The mixture was stirred at room temperature for a further 16 hr. The solvent was removed *in vacuo* and the crude material was purified by flash chromatography on silica (eluant methanol:dichloromethane 5:95); yield 0.115g (82%).

M.S.: calc $m/e=462.26$; found $[M+H]^+=463.2$

Example 40

(4R)-4-Hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-N,N-dimethylamide



40A. L-trans-4-Hydroxyproline-N,N-dimethylamide hydrochloride

To a solution of BOC-hydroxyproline (2.99g, 13.89mmol) in dichloromethane (100ml) were added *N,N*-diisopropylethylamine (3.7ml, 21.24mmol), 4-(dimethylamino)pyridine (1.74g, 14.24mmol), dimethylamine hydrochloride (1.72g, 21.09mmol) and WSCDI (3.17g, 16.68mmol). The mixture was stirred at room temperature for 30h. The mixture was diluted with dichloromethane (100ml) and washed with 0.3M KHSO_4 , saturated sodium bicarbonate solution and brine, dried over MgSO_4 , and concentrated *in vacuo* to give a colourless gum. This crude material was taken up in 4N HCl/dioxan (50ml) and stirred at room temperature for 1hr and then concentrated *in vacuo*. The residue was azeotroped with toluene and diethyl ether to give a white solid; yield 0.45g (17%).


40B. (4R)-4-Hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzyl-carbamoyl)-L-proline-N,N-dimethylamide.

The amine of Example 39B (0.10g, 0.302mmol) was reacted with the amine of Example 40A (0.153mg, 0.785mmol) following the method of Example 39C. The product was purified by flash chromatography on silica (eluant chloroform:methanol:acetic acid 95:4:1); yield 0.95g (66%).

M.S.: calc m/e=478.26; found $[M+H]^+ = 479.2$


Following the above methods, the following compounds were also prepared.

Table A – Examples 41 - 44




Ex.	Ar	W	Z	M+H ⁺
41	Ph	N	CH=CH	465.4
42	Ph	CH	S	470.2
43	4-Me-Ph	CH	CH=CH	478.1
44	2-Me-Ph	CH	CH=CH	478.1


Table B – Examples 45 - 55



Ex.	Ar	R ¹	R ²	R ³	A ¹⁶	V	M+H ⁺
45	2,6-F ₂ -Ph	H	H	H	NCH ₂ Ph	O	527.4
46	2,6-F ₂ -Ph	H	H	H	S	O	454
47	1-Nap	H	H	Cl	CH ₂	O	484
48	Ph	H	H	Cl	CH ₂	O	434
49	3-Pyr	H	H	Me	CH ₂	S	431.1
50	2,6-F ₂ -Ph	MeO	H	H	CH ₂	O	466
51	2,6-F ₂ -Ph	H	CH:CH-CH:CH		CH ₂	O	486
52	2,6-F ₂ -Ph	H	H	Me	N(CH ₂) ₂ NMe ₂	O	522.3
53	2,6-F ₂ -Ph	Cl	H	Cl	CH ₂	O	504.1
54	2,6-F ₂ -Ph	H	H	Me	SO ₂	O	500.2
55	2,6-F ₂ -Ph	H	H	Me	NCH ₂ CO ₂ H	O	509.2

Table C – Examples 56 - 57



Ex.	G ¹	M+H ⁺
56		488.3



Ex.	G ¹	M+H ⁺
57		517.1

Table D – Examples 58 - 61









Ex.	G ¹	R ⁸	M+H ⁺
58		OEt	439
59		NMe ₂	473.3
60		NMe ₂	461.1
61		NMe ₂	476

Table E – Examples 62 - 70



Ex.	f	R ²	R ³	R ⁸	M+H ⁺
62	1	H	Me		463
63	1	H	Me		449.2
64	0	H	Me	OE _t	410
65	1	Me	H	OE _t	424
66	1	H	Me	O _i Pr	438
67	1	H	Me	O _i Bu	452
68	1	H	Cl	NMe ₂	443
69	2	H	Me	OE _t	438
70	2	H	Me	OH	410


Table F – Examples 71 - 77



Ex.	A ¹⁶	f	R ²	R ³	R ⁸	M+H ⁺
71	O	2	H	H	OMe	397

Ex.	A ¹⁸	f	R ²	R ³	R ⁸	M+H ⁺
72	CH ₂	1	H	Me	OMe	415
73	CH ₂	1	H	Me	OEt	409
74	CH ₂	1	H	Me	OH	381
75	CH ₂	2	H	Me	OH	395
76	CH ₂	3	H	Me	OH	409
77	CH ₂	1	Me	H	OMe	395

Table G – Examples 78 - 90.



Ex.	G ¹	R ³	M+H ⁺
78		Me	502
79		OMe	479.2
80		Et	477.3
81		Me	479.2
82		Me	518.0










Ex.	G ¹	R ³	M+H ⁺
83		Me	532.2
84		Me	517.2
85		Me	513.7
86		Me	527.0
87		Me	514.6
88		Me	516.1
89		Me	515.0
90		Me	500.7

Table H – Examples 91 - 106



Ex.	G ¹	E ¹	E ²	M+H ⁺
91		H	OAc	521.0
92		=O		477.3
93		H	OBn	638.2
94		Br	H	541.1
95		F	F	499.2
96		H	OBn	619.2
97		H	N ₃	504.3
98		H	O-tBu	535.3
99		H	OH	517.6









Ex.	G ¹	E ¹	E ²	M+H ⁺
100		H	OH	546.3
101		H	OH	547.9
102		H	OMe	548.2
103		H	OMe	562.1
104		H	Cl	566.2
105		H	NHBn	568
106		OCH ₂ CH ₂ O		558.3

Table I – Examples 107 - 124



Ex.	G ¹	R ³	F ¹	F ²	L	M+H ⁺
107		Me	=O		NMe ₂	493.5
108		Me	=O		NMe ₂	530.3
109		Me	=O		NMe ₂	543.4
110		Me	=O		NMe ₂	532.4
111		Me	=O		NMe ₂	544.3
112		Me	=O		NMe ₂	536.4
113		Me	=O		NMe ₂	494.5
114		Cl	=O		NMe ₂	515.2













Ex.	G ¹	R ³	F ¹	F ²	L	M+H ⁺
115		Cl	=O		NMe ₂	551.5
116		Me	=O		NMeEt	558.3
117		Me	=O			570.3
118		Me	=S		NMe ₂	546.2
119		Cl	=S		NMe ₂	535.1
120		Cl	=S		NMe ₂	585.1
121		Me	=O		NMe ₂	590.2
122		Me	=O		NMe ₂	548.2
123		Me	=O		NMe ₂	494.3
124		Me	=O		NMe ₂	522.4


Table J – Examples 125 - 153



Ex.	E ¹	E ²	F ¹	F ²	L		M+H ⁺
125	H	H	H	H	OMe	R	436.4
126	H	H	H	H	OMe	S	436.2
127	H	H	=O		NMeEt	R	477.2
128	H	OPh	=O		OMe	R	542.3
129	H	OPh	=O		OH	RS	528.3
130	H	OPh	=O		NMe ₂	RS	555.3
131	H	F	=O		OH	R	454.4
132	OMe	OMe	=O		OMe	R	510.3
133	OMe	OMe	=O		OH	R	496.2
134	H	H	=O		OBu	R	492.5
135	H	H	=O		OH	R	436.3
136	H	OH	=O		OMe	R	466.0
137	H	OH	=O		OEt	R	480.2
138	H	H	=S		NMe ₂	R	479.2
139	H	OMe	=O		OMe	R	480.2
140	H	H	=O		OiPr	R	478.2
141	H	OH	=O		OH	R	452.1
142	H	OBn	=O		OiPr	R	584.2
143	H	OH	=O		OiPr	R	494.1
144	H	OBn	=O		NMe ₂	R	569.2
145	H	OMe	=O		OH	R	466.2
146	H	OEt	=O		NMe ₂	R	507.3
147	H	Cl	=O		OMe	R	484.1


Ex.	E ¹	E ²	F ¹	F ²	L		M+H ⁺
148	H	Cl	=O		OH	R	470.1
149	H	Cl	=O		NMe ₂	R	497.2
150	Cl	H	=O		NMe ₂	R	497.2
151	H	F	=O		OMe	R	468.3
152	H	F	=O		NMe ₂	R	481.3
153	OMe	OMe	=O		NMe ₂	R	523.3

Table K – Examples 154 - 159




Ex.	R ²	R ³	E ²	F ¹	F ²	M+H ⁺
154	H	Cl	H	=O		483.4
155	Me	H	H	=O		463.2
156	Cl	H	H	=O		483.1
157	H	Cl	H	=S		499.2
158	H	Cl	OBn	=O		589.2
159	H	Cl	OH	=O		499.2

Table L – Examples 160 - 164




Ex.	R ²	E ²	M+H ⁺
160	Cl	H	489.1
161	Me	H	469.2
162	Me	OH	485.0
163	Cl	OMe	519.3
164	Me	OMe	499.3

Table M – Examples 165 - 170




Ex.	R ⁴	E ²	F ¹	F ²	V	M+H ⁺
165	H	H	=O		S	479.4
166	H	OH	=O		S	495.0
167	H	H	=S		S	495.1
168	Me	H	=O		O	477.2
169	H	OBn	=O		S	585.2
170	H	OBn	=S		O	585.0

Table N – Examples 171 - 177



Ex.	E ¹	E ²	F ¹	F ²	L	M+H ⁺
171	H	H	=S		NMe ₂	516.2
172	H	OBn	=O		NMe ₂	606.3
173	H	OH	=O		NMe ₂	507.3
174	OMe	OMe	=O		OMe	547.3
175	-OCH ₂ CH ₂ O-		=O		OMe	545.3
176	-OCH ₂ CH ₂ O-		=O		NMe ₂	558.3
177	-SCH ₂ CH ₂ S-		=O		NMe ₂	590.2


Table O— Examples 178 - 182



Ex.	E ¹	E ²	F ¹	F ²	L	M+H ⁺
178	H	OH	=O		NMe ₂	516.1

Ex.	E ¹	E ²	F ¹	F ²	L	M+H ⁺
179	H	H	=S		NMe ₂	516.2
180	H	OMe	=O		NMe ₂	530.4
181	-OCH ₂ CH ₂ O-		=O		OMe	545.3
182	-OCH ₂ CH ₂ O-		=O		OH	531.3

Table P- Examples 183 - 190



Ex.	A ¹⁰	R ³	E ²	F ¹	F ²	M+H ⁺
183	O	Me	H	=O		519.3
184	NMe	Me	H	=O		532.33
185	NMe	Me	OH	=O		548.1
186	NMe	Me	OMe	=O		562.3
187	O	Me	OMe	=O		549.2
188	NMe	Me	OMe	=S		578.2
189	O	Cl	OMe	=O		569.1
190	O	Me	OMe	=S		565.2

Table Q – Representative NMR data

Ex. No	¹ H NMR (CDCl ₃)
28	δ 1.40-1.60 (1H, m), 1.84-2.20 (3H, m), 2.15 (3H, s), 2.40-2.54 (2H, m), 2.58-2.92 (4H, m), 2.94-3.10 (1H, m), 3.65 (3H, s), 4.30 (2H, d, J=5.6Hz), 4.99 (1H, d, J=12.9Hz), 5.90 (1H, s), 6.62 (1H, d, J=7.9Hz), 6.78-6.96

	(3H, m), 7.00-7.16 (2H, m), 7.21 (1H, m) ppm
29	δ 1.48-1.70 (1H, m), 1.96-2.16 (3H, m), 2.26 (3H, s), 2.78-3.18 (3H, m), 3.98 (3H, s), 4.50 (2H, d, J=6.8Hz), 5.08 (1H, d, J=12.7Hz), 6.72 (1H, d, 7.6Hz), 6.88-7.06 (3H, m), 7.18 (1H, t, J=7.6Hz), 7.22-7.36 (2H, m) ppm
30	δ 1.40-1.62 (1H, m), 1.84-2.24 (3H, s), 2.17 (3H, s), 2.70-3.10 (3H, m), 4.40 (2H, d, J=5.9Hz), 4.99 (1H, d, J=12.9Hz), 6.63 (1H, d, J=7.6Hz), 6.80-6.98 (3H, m), 7.02-7.28 (3H, m), 7.38 (1H, br s) ppm
31	δ 1.42-1.62 (1H, m), 1.84-2.28 (8H, m), 2.30-2.50 (4H, m), 2.70-2.94 (2H, m), 2.96-3.12 (1H, m), 3.65 (3H, s), 4.31 (2H, d, J=5.3Hz), 4.99 (1H, d, J=13.9Hz), 5.75 (1H, br s), 6.63 (1H, d, J=7.6Hz), 6.78-6.98 (3H, m), 7.02-7.16 (2H, m), 7.21 (1H, d, J=6.6Hz) ppm
32	δ 1.18 (3H, t, J=7.3Hz), 1.38-1.55 (1H, m), 1.80-2.10 (3H, m), 1.95 (3H, s), 2.60-2.98 (3H, m), 3.84 (2H, s), 4.04 (2H, s), 4.07 (2H, q, J=7.3Hz), 4.87-4.92 (1H, m), 5.73 (2H, br s), 6.50 (1H, d, J=7.3Hz), 6.63-6.97 (5H, m), 7.11 (1H, d, J=7.3Hz) ppm
33	δ 1.30-1.50 (1H, m), 1.75-2.05 (3H, m), 1.94 (3H, s), 2.60-2.98 (3H, m), 3.59 (2H, br s), 4.01 (2H, br s), 4.80-4.85 (1H, m), 6.05 (2H, br s), 6.53 (1H, d, J=7.2Hz), 6.75-6.99 (5H, m), 7.11 (1H, d, J=7.2Hz) ppm.
34	δ 1.40-1.60 (1H, m), 1.80-2.00 (2H, m), 2.00-2.20 (3H, s), 2.60 (3H, d, J=4.0Hz), 2.65-3.05 (3H, m), 3.60 (2H, d, J=4.0Hz), 4.15 (2H, d, J=4.0Hz), 4.90-5.00 (1H, m), 6.10-6.30 (2H, m), 6.60 (1H, d, J=8.0Hz), 6.70-7.20 (8H, m) ppm
35	δ 1.39-1.50 (1H, m), 1.86-2.10 (3H, m), 2.07 (3H, s), 2.57 (3H, s), 2.60-3.00 (3H, m), 2.85 (3H, s), 3.95 (2H, d, J=4.0Hz), 4.16 (2H, d, J=5.6Hz), 4.90-5.00 (1H, m), 5.74 (1H, br s), 6.11 (1H, br s), 6.54 (1H, d, J=7.6Hz), 6.78-7.18 (6H, m) ppm
36	δ 1.38-1.50 (1H, m), 1.80-2.00 (3H, m), 2.00 (3H, s), 2.60-3.00 (3H, m), 3.64 (3H, s), 3.90 (2H, s), 4.10 (2H, s), 4.85-4.95 (1H, m), 6.52 (1H, d, J=7.2Hz), 6.67-7.02 (7H, m), 7.13 (1H, d, J=6.2Hz) ppm
37	δ 1.40-1.76 (2H, m), 1.84-2.16 (2H, m), 2.29 (3H, s), 2.66-3.10 (3H, s), 3.95 (2H, s), 4.56 (2H, s), 4.99 (1H, d, J=13.9Hz), 5.59 (1H, br s), 6.63 (1H, d, J=7.9Hz), 6.80-6.98 (3H, m), 7.00-7.12 (2H, m), 7.20 (1H, d, J=7.3Hz) ppm
38	δ 1.70-1.86 (3H, m), 1.96-2.08 (2H, m), 2.44-2.56 (2H, m), 2.60-2.72 (2H,

	m), 2.86-2.98 (2H, m), 3.67 (3H, s), 3.85 (1H, br s), 4.44 (2H, d, J=5.9Hz), 6.18 (1H, d, J=5.3Hz), 6.28 (1H, br s), 6.68 (1H, d, J=5.3Hz), 7.03 (1H, d, J=7.6Hz), 7.15 (1H, d, J=7.6Hz) ppm
39	δ 1.35-1.55 (1H, m), 1.74-2.10 (3H, m), 2.11 (3H, s), 2.17-2.35 (1H, m), 2.60-2.82 (2H, m), 2.86 (3H, s), 2.90-3.14 (2H, m), 3.05 (3H, s), 3.26 (1H, dd, J=14.9 & 7.2Hz), 3.40-3.53 (1H, m), 3.64-3.84 (1H, m), 4.03-4.19 (1H, m), 4.29-4.42 (1H, m), 4.55-4.68 (1H, m), 4.74-4.81 (1H, m), 4.85-4.98 (1H, m), 6.58 (1H, d, J=7.7Hz), 6.75-6.89 (2H, m), 6.91-7.06 (3H, m), 7.16 (1H, d, J=6.5Hz), 7.93-8.03 (1H, m) ppm
40	δ 1.65-1.80 (2H, m), 1.85-2.00 (3H, m), 2.05-2.25 (1H, m), 2.10 (3H, s), 2.80-3.10 (3H, m), 2.85 (3H, s), 3.00 (3H, s), 3.40-3.30 (1H, m), 3.45-3.55 (1H, m), 3.65-3.95 (1H, m), 4.00-4.10 (1H, m), 4.30-4.55 (1H, m), 4.91 (1H, t, J=7.7Hz), 5.15-5.30 (1H, m), 6.10-6.20 (1H, m), 6.55-6.65 (1H, m), 6.85-7.50 (5H, m) ppm
66	δ 1.17 (6H, d, J=6.3Hz), 1.20-1.24 (1H, m), 1.80-2.10 (3H, m), 2.00 (3H, s), 2.60-3.00 (3H, m), 3.85 (2H, d, J=5.3Hz), 4.10 (2H, d, J=4.9Hz), 4.82-4.85 (1H, m), 4.96 (1H, sept, J=6.2Hz), 5.33 (1H, t, J=5.2Hz), 5.43 (1H, t, J=4.9Hz), 6.52 (1H, d, J=7.6Hz) ppm
67	δ 1.38-1.42 (1H, m), 1.38 (9H, s), 1.78-2.10 (3H, m), 1.97 (3H, s), 2.60-3.00 (3H, m), 3.78 (2H, s), 4.07 (2H, s), 4.89-4.94 (1H, m), 5.50 (2H, br s), 6.51 (1H, d, J=7.9Hz), 6.64-6.98 (5H, m), 7.12 (1H, d, J=7.7Hz) ppm
68	δ 1.38-1.50 (1H, m), 1.80-2.06 (3H, m), 2.60-3.00 (3H, m), 2.70 (3H, s), 2.87 (3H, s), 3.96 (2H, d, J=4.0Hz), 4.27 (2H, d, J=6.0Hz), 4.85-4.95 (1H, m), 5.98 (1H, t, J=6.0Hz), 6.14 (1H, t, J=4.0Hz), 6.55 (1H, d, J=7.6Hz), 6.80-7.16 (6H, m) ppm
69	δ 1.25 (3H, t, J=7.0Hz), 1.40-1.60 (1H, m), 1.85-2.20 (3H, m), 2.04 (3H, s), 2.45 (2H, t, J=6.27Hz), 2.65-3.10 (3H, m), 3.30-3.50 (2H, m), 4.00-4.20 (4H, m), 4.90-5.00 (1H, m), 5.50-5.70 (2H, m), 6.50-7.20 (7H, m) ppm
70	δ 1.20-1.45 (1H, m), 1.65-2.05 (3H, m), 1.95 (3H, s), 2.05-2.25 (2H, m), 2.50-3.00 (3H, m), 3.00-3.20 (2H, m), 3.85-4.05 (2H, m), 4.65-4.90 (1H, m), 5.80-6.20 (1H, brs), 6.40-7.20 (9H, m) ppm

Example 191.**Determination of V₂ receptor agonist activity *in vitro***

Agonist activity was determined for all compounds and is reported as an EC₅₀ value, being that concentration of compound necessary to cause a half-maximal cellular activation. All the compounds had EC₅₀ values of 10 μ M or less, and typical results are listed in Table R.

Table R – EC₅₀ values for typical compounds

Compound of Example	EC ₅₀ (nM)	Compound of Example	EC ₅₀ (nM)
1	39	15	44
2	160	16	16
3	300	17	16
4	300	18	17
5	150	19	40
6	47	20	17
7	24	21	180
8	220	22	1000
9	50	23	40
10	4	24	92
11	21	25	280
12	50	26	10
13	38	27	23
14	240		

Example 192.**Determination of antidiuretic activity *in vivo***

The Brattleboro rat is a recognised model for vasopressin deficiency (for a review see FD Grant, "Genetic models of vasopressin deficiency", *Exp. Physiol.* **85**, 203S-209S, 2000). The animals do not secrete vasopressin and consequently produce large volumes of dilute urine. Compounds of the invention were administered to Brattleboro rats (0.1-10mg/kg p.o. in methylcellulose). Urine was collected hourly and volumes were compared with control animals. Animals had free access to food and water throughout the experiment. Representative results are given in Table S. Results for Desmopressin are given for comparison.

Table S – Antidiuretic activity

Compound of Example	Dose	% inhibition of urine output (at 1 hour)
32	1mg/kg	74
33	1mg/kg	38
35	1mg/kg	45-82
39	1mg/kg	82
62	1mg/kg	58
88	1mg/kg	60
103	1mg/kg	63
107	1mg/kg	84
119	1mg/kg	68
163	1mg/kg	90
Desmopressin	0.1mg/kg	37
	1mg/kg	100
	10mg/kg	100

Example 193.**Pharmaceutical composition for tablet**

Tablets containing 100mg of the compound of Example 39 as the active agent are prepared from the following:

Compound of Example 39	200.0g
Corn starch	71.0g
Hydroxypropylcellulose	18.0g
Carboxymethylcellulose calcium	13.0g
Magnesium stearate	3.0g
Lactose	195.0g
<i>Total</i>	<i>500.0g</i>

The materials are blended and then pressed to give 2000 tablets of 250mg, each containing 100mg of the compound of Example 39.


The foregoing Examples demonstrate that compounds within the scope of the invention are readily prepared using standard chemical techniques, and that these compounds have the biological properties that would be expected of V_2 receptor agonists. In particular, the compounds are potent antidiuretics in an animal model of vasopressin deficiency. Thus it is clear that they may be useful in the treatment of human diseases that are currently treatable with Desmopressin, such as central diabetes insipidus, nocturnal enuresis and nocturia. It has further been suggested that antidiuretics such as Desmopressin may be useful in certain types of urinary incontinence. These arguments would also extend to the compounds of the present invention.

Desmopressin is also used in the treatment of certain coagulation disorders. There is good evidence to suggest that this action is also mediated through the V_2 receptor (see for example JE Kaufmann *et al.*, "Vasopressin-induced von Willebrand factor secretion from endothelial cells involves V_2 receptors and cAMP", *J. Clin. Invest.* **106**, 107-116, 2000; A Bernat *et al.*, " V_2 receptor antagonism of DDAVP-induced release of hemostasis factors in conscious dogs", *J. Pharmacol. Exp. Ther.* **282**, 597-602, 1997), and hence it would be expected that the compounds of the present invention should be useful pro-coagulants.

The scope of the present invention is further defined in the following Claims.

CLAIMS

- 1 A compound according to general formula 1 or 2, or a tautomer or a pharmaceutically acceptable salt thereof,




wherein:

W is either N or C-R⁴;

R¹ – R⁴ are independently selected from H, F, Cl, Br, alkyl, CF₃, phenyl, OH, O-alkyl, NH₂, NH-alkyl, N(alkyl)₂, NO₂ and CN, or R² and R³ together can be –CH=CH-CH=CH–;

G¹ is a bicyclic or tricyclic fused azepine derivative selected from general formulae 3 to 8,



in which A¹, A⁴, A⁷ and A¹⁰ are each independently selected from CH₂, O and NR⁵;

A², A³, A⁹, A¹¹, A¹³, A¹⁴ and A¹⁵ are each independently selected from CH and N;

either A⁵ is a covalent bond and A⁶ is S, or A⁵ is N=CH and A⁶ is a covalent bond;

A⁸ and A¹² are each independently selected from NH, NCH₃ and S;

A¹⁶ and A¹⁷ are both CH₂, or one of A¹⁶ and A¹⁷ is CH₂ and the other is selected from

CH(OH), CF₂, O, SO_a and NR⁶;

R⁵ is selected from H, alkyl, CO-alkyl and (CH₂)_bR⁶;

R⁶ is selected from phenyl, pyridyl, OH, O-alkyl, NH₂, NH-alkyl, N(alkyl)₂, NO₂, CO₂H and CN;


a is 0, 1 or 2;

b is 1, 2, 3 or 4;


Y is CH or N;

Z is CH=CH or S; and


G² is a group selected from general formulae 9 to 11,



9



10



11

in which Ar is selected from phenyl, pyridyl, naphthyl and mono- or polysubstituted phenyl or pyridyl wherein the substituents are selected from F, Cl, Br, alkyl, OH, O-alkyl, NH₂, NH-alkyl, N(alkyl)₂, NO₂ and CN;

D is a covalent bond or NH;

E¹ and E² are both H, OMe or F, or one of E¹ and E² is OH, O-alkyl, OBn, OPh, OAc, F, Cl, Br, N₃, NH₂, NHBn or NHAc and the other is H, or E¹ and E² together are =O, -O(CH₂)₉O- or -S(CH₂)₉S-;

F¹ and F² are both H, or together are =O or =S;

L is selected from OH, O-alkyl, NH₂, NH-alkyl and NR⁹R¹⁰;

R⁷ is selected from H, alkyl, alkenyl and COR⁸;

R⁸ is selected from OH, O-alkyl, NH₂, NH-alkyl, N(alkyl)₂, pyrrolidinyl and piperidinyl;

R⁹ and R¹⁰ are both alkyl, or together are -(CH₂)_h- or -(CH₂)₂O(CH₂)_z;

V is O, N-CN or S;

c is 0 or 1;

d is 0 or 1;

e is 0 or 1;

f is 0, 1, 2, 3 or 4;

g is 2 or 3; and

h is 3, 4 or 5,

provided that d and e are not both 0.

- 2 A compound according to Claim 1, or a tautomer or pharmaceutically acceptable salt thereof, wherein the compound is a compound according to general formula 1.
- 3 A compound according to Claim 2, or a tautomer or pharmaceutically acceptable salt thereof, wherein W is C-R⁴.
- 4 A compound according to Claim 3, or a tautomer or pharmaceutically acceptable salt thereof, wherein at least one of R¹ – R⁴ is not H.
- 5 A compound according to Claim 4, or a tautomer or pharmaceutically acceptable salt thereof, wherein one of R¹ – R⁴ is methyl, F or Cl and the others are all H.
- 6 A compound according to Claim 1, or a tautomer or pharmaceutically acceptable salt thereof, wherein the compound is a compound according to general formula 2.
- 7 A compound according to any preceding Claim, or a tautomer or pharmaceutically acceptable salt thereof, wherein G¹ is a group according to one of general formulae 3 to 7.
- 8 A compound according to Claim 7, or a tautomer or pharmaceutically acceptable salt thereof, wherein Y is CH.
- 9 A compound according to Claim 8, or a tautomer or pharmaceutically acceptable salt thereof, wherein Z is –CH=CH–.
- 10 A compound according to Claim 8, or a tautomer or pharmaceutically acceptable salt thereof, wherein Z is S.
- 11 A compound according to Claim 7, or a tautomer or pharmaceutically acceptable salt thereof, wherein Y is N and Z is –CH=CH–.
- 12 A compound according to any of Claims 7 to 11, or a tautomer or pharmaceutically acceptable salt thereof, wherein G¹ is a group according to general formula 3.

- 13 A compound according to Claim 12, or a tautomer or pharmaceutically acceptable salt thereof, wherein A^1 is CH_2 and A^2 and A^3 are both CH.
- 14 A compound according to any of Claims 7 to 11, or a tautomer or pharmaceutically acceptable salt thereof, wherein G^1 is a group according to general formula 6.
- 15 A compound according to Claim 14, or a tautomer or pharmaceutically acceptable salt thereof, wherein A^{11} is CH and A^{12} is S.
- 16 A compound according to any of Claims 1 to 6, or a tautomer or pharmaceutically acceptable salt thereof, wherein G^1 is a group according to general formula 8.
- 17 A compound according to Claim 16, or a tautomer or pharmaceutically acceptable salt thereof, wherein A^{17} is CH_2 .
- 18 A compound according to either Claim 16 or 17, or a tautomer or pharmaceutically acceptable salt thereof, wherein A^{16} is CH_2 .
- 19 A compound according to any preceding Claim, or a tautomer or pharmaceutically acceptable salt thereof, wherein G^2 is a group according to general formula 9.
- 20 A compound according to Claim 19, or a tautomer or pharmaceutically acceptable salt thereof, wherein Ar is mono- or polysubstituted phenyl.
- 21 A compound according to either Claim 19 or 20, or a tautomer or pharmaceutically acceptable salt thereof, wherein Ar is phenyl substituted with at least two halogen atoms selected from F and Cl.
- 22 A compound according to any of Claims 19 to 21, or a tautomer or pharmaceutically acceptable salt thereof, wherein Ar is 2,6-difluorophenyl.
- 23 A compound according to any of Claims 1 to 18, or a tautomer or pharmaceutically acceptable salt thereof, wherein G^2 is a group according to general formula 10.
- 24 A compound according to Claim 23, or a tautomer or pharmaceutically acceptable salt

thereof, wherein R^7 is COR^8 .

- 25 A compound according to Claim 24, or a tautomer or pharmaceutically acceptable salt thereof, wherein R^8 is $N(alkyl)_2$.
- 26 A compound according to any of Claims 1 to 18, or a tautomer or pharmaceutically acceptable salt thereof, wherein G^2 is a group according to general formula 11.
- 27 A compound according to Claim 26, or a tautomer or pharmaceutically acceptable salt thereof, wherein F^1 and F^2 together are $=O$.
- 28 A compound according to either Claim 26 or 27, or a tautomer or pharmaceutically acceptable salt thereof, wherein E^1 and E^2 are both H or one is H and the other is O-alkyl.
- 29 A compound according to any of Claims 26 to 28, or a tautomer or pharmaceutically acceptable salt thereof, wherein one of E^1 and E^2 is H and the other is O-alkyl, and the stereochemistry at the CE^1E^2 centre is of the *R* absolute configuration.
- 30 A compound according to any of Claims 26 to 29, or a tautomer or pharmaceutically acceptable salt thereof, wherein the stereochemistry adjacent to the ring nitrogen is of the *S* absolute configuration.
- 31 A compound according to Claim 1, or a tautomer or pharmaceutically acceptable salt thereof, selected from

1-(4-[3-(2-Chloro-6-fluorophenyl)ureidomethyl]-3-methylbenzoyl)-2,3,4,5-tetrahydro-1*H*-1-benzazepine,

1-(4-[3-(2,6-Difluorophenyl)ureidomethyl]-3-methylbenzoyl)-5-(3-pyridyl)methyl-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepine,

1-(3-Chloro-4-[3-(2-chloro-6-fluorophenyl)ureidomethyl]benzoyl)-5-ethyl-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepine,

4-(3-Chloro-4-[3-(2,6-difluorophenyl)ureidomethyl]benzoyl)-5,6,7,8-tetrahydrothieno[3,2-*b*]azepine,

1-(3-Chloro-4-(3-(methyloxycarbonyl)propanoylaminomethyl)benzoyl)-2,3,4,5-tetrahydro-1-benzazepine,

1-(2-Methyl-4-(5-(3-pyridylmethyl)-2,3,4,5-tetrahydro-1,5-benzodiazepin-1-ylcarbonyl)benzyl)-3-(methyloxycarbonylmethyl)urea,

1-(2-Methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-Hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(3-Chloro-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(2-Chloro-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-4-Benzyloxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-Methoxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-Methoxy-1-(3-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(2-Chloro-4-(5,6,7,8-tetrahydro-4*H*-thieno[3,2-*b*]azepin-4-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(4-(10,11-Dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-yl carbonyl)-2-methylbenzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(2-Chloro-4-(10,11-Dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)-benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide, and

(4*R*)-1-(4-(10,11-Dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)-2-methylbenzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylthioamide.

- 32 A compound according to Claim 1, or a tautomer or pharmaceutically acceptable salt thereof, selected from

1-(2-Methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide, and

(4*R*)-4-Hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide.

- 33 A use for a compound according to any of Claims 1 to 32 or a pharmaceutically acceptable salt thereof, which use is as a component of a pharmaceutical composition.
- 34 A use for a compound according to any of Claims 1 to 32 or a pharmaceutically acceptable salt thereof, which use is as a therapeutic agent for the treatment of nocturnal enuresis, nocturia, polyuria resulting from central diabetes insipidus, urinary incontinence or bleeding disorders.
- 35 A pharmaceutical composition which comprises, as an active agent, a compound according to any of Claims 1 to 32.
- 36 A pharmaceutical composition according to Claim 35, which composition is to be used for the treatment of polyuria.
- 37 A pharmaceutical composition according to Claim 35, which composition is to be used for the control of urinary incontinence.
- 38 A pharmaceutical composition according to Claim 37, which composition is for voiding postponement.

- 39 A pharmaceutical composition according to Claim 35, which composition is to be used for the treatment of bleeding disorders.
- 40 A method of treatment of nocturnal enuresis, nocturia and diabetes insipidus, which method comprises the administration to a person in need of such treatment of an effective amount of a composition according to Claim 35.
- 41 A method for the control of urinary incontinence, which method comprises the administration to a person in need of such treatment of an effective amount of a composition according to Claim 35.
- 42 A method for the control of urinary incontinence according to Claim 41, wherein the treatment results in voiding postponement.
- 43 A method for the treatment of bleeding disorders, which method comprises the administration to a person in need of such treatment of an effective amount of a composition according to Claim 35.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 01/02737

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D223/16 A61K31/55 A61P13/00 C07D401/06 C07D243/12
C07D495/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	WO 99 06409 A (AMERICAN HOME PROD) 11 February 1999 (1999-02-11) cited in the application claim 1 ---	1-13, 23-25, 33-43 14-22, 26-32
A	WO 99 06403 A (AMERICAN HOME PROD) 11 February 1999 (1999-02-11) cited in the application claim 1 ---	1-43
A	WO 95 34540 A (TOMINAGA MICHIAKI ;YAMASHITA HIROSHI (JP); KAN KEIZO (JP); KONDO K) 21 December 1995 (1995-12-21) page 8, line 3 -page 9, line 10; claim 1 --- -/--	1-43

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search

17 September 2001

Date of mailing of the international search report

24/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Uselli, A

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/GB 01/02737

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	GB 2 355 454 A (FERRING BV) 25 April 2001 (2001-04-25) page 1 -page 3; claim 1 ---	1-43
E	WO 01 49682 A (FRANKLIN RICHARD JEREMY ;HUDSON PETER (GB); ASHWORTH DOREEN MARY () 12 July 2001 (2001-07-12) page 3, line 1 -page 4, line 17; claim 1 -----	1-43

International Application No

100/GB 01/02737

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9906409 A	11-02-1999	AU 8663398 A	22-02-1999
		BR 9811585 A	26-09-2000
		CN 1272111 T	01-11-2000
		EP 1000062 A	17-05-2000
		HU 0002480 A	28-11-2000
		NO 20000242 A	06-03-2000
WO 9906403 A	11-02-1999	AU 8592598 A	22-02-1999
		BR 9811559 A	12-09-2000
		CN 1272110 T	01-11-2000
		EP 1000059 A	17-05-2000
		HU 0003394 A	28-08-2001
		NO 20000241 A	06-03-2000
WO 9534540 A	21-12-1995	AU 690283 B	23-04-1998
		AU 2629395 A	05-01-1996
		CA 2192928 A	21-12-1995
		CN 1150799 A	28-05-1997
		EP 0765314 A	02-04-1997
		JP 11349570 A	21-12-1999
		JP 8301848 A	19-11-1996
		JP 2000351768 A	19-12-2000
		US 6096735 A	01-08-2000
GB 2355454 A	25-04-2001	AU 1037401 A	30-04-2001
		WO 0129005 A	26-04-2001
WO 0149682 A	12-07-2001	NONE	